

**Solutions of Sample Problems for First In-Class Exam
Math 246, Fall 2008, Professor David Levermore**

- (1) (a) Write a MATLAB command that evaluates the definite integral

$$\int_0^{\infty} \frac{r}{1+r^4} dr.$$

Solution: The simplest solution is

$$\text{int('x/(1+x^4)', 'x', 0, inf),}$$

where you can replace x by any other letter or use `Inf` instead of `inf`.

- (b) Sketch the graph that you expect would be produced by the following MATLAB commands.

```
[x, y] = meshgrid(-5:0.5:5, -5:0.2:5)
contour(x, y, x.^2 + y.^2, [25, 25])
axis square
```

Solution: Your sketch should show both x and y axes marked from -5 to 5 and a single circle of radius 5 centered at the origin. The tick marks on the axes should mark intervals of length $.5$.

- (2) Find the explicit solution for each of the following initial-value problems and identify its interval of existence (definition).

(a) $\frac{dz}{dt} = \frac{\cos(t) - z}{1+t}, \quad z(0) = 2.$

Solution: This equation is *linear* in z , so write it in the linear normal form

$$\frac{dz}{dt} + \frac{z}{1+t} = \frac{\cos(t)}{1+t}.$$

An integrating factor is given by

$$\exp\left(\int_0^t \frac{1}{1+s} ds\right) = \exp(\log(1+t)) = 1+t,$$

Upon multiplying the equation by $(1+t)$, one finds that

$$\frac{d}{dt}((1+t)z) = \cos(t),$$

which is then integrated to obtain

$$(1+t)z = \sin(t) + C.$$

The integration constant C is found through the initial condition $z(0) = 2$ by setting $t = 0$ and $z = 0$, whereby

$$C = (1+0)2 - \sin(0) = 2.$$

Hence, upon solving explicitly for z , the solution is

$$z = \frac{2 + \sin(t)}{1+t}.$$

The interval of existence for this solution is $t > -1$.

(b) $\frac{du}{dz} = e^u + 1, \quad u(0) = 0.$

Solution: This equation is *autonomous* (and therefore *separable*), so write it in the separated differential form

$$\frac{1}{e^u + 1} du = dz.$$

This equation can be integrated to obtain

$$z = \int \frac{1}{e^u + 1} du = \int \frac{e^{-u}}{1 + e^{-u}} du = -\log(1 + e^{-u}) + C.$$

The integration constant C is found through the initial condition $u(0) = 0$ by setting $z = 0$ and $u = 0$, whereby

$$C = 0 + \log(1 + e^0) = \log(2).$$

Hence, the solution is given implicitly by

$$z = -\log(1 + e^{-u}) + \log(2) = -\log\left(\frac{1 + e^{-u}}{2}\right).$$

This may be solve for u as follows:

$$\begin{aligned} e^{-z} &= \frac{1 + e^{-u}}{2}, \\ 2e^{-z} - 1 &= e^{-u}, \\ u &= -\log(2e^{-z} - 1). \end{aligned}$$

The interval of existence for this solution is $z < \log(2)$.

(3) Consider the differential equation

$$\frac{dy}{dt} = 4y^2 - y^4.$$

(a) Find all of its stationary (equilibrium) solutions and classify each as being either stable, unstable, or semistable.


Solution: The right-hand side of the equation factors as

$$4y^2 - y^4 = y^2(4 - y^2) = y^2(2 + y)(2 - y),$$

which implies that $y = -2$, $y = 0$, and $y = 2$ are all of its stationary solutions. A sign analysis of $y^2(2 + y)(2 - y)$ then shows that

$$\begin{aligned} \frac{dy}{dt} &> 0 && \text{when } -2 < y < 0 \text{ or } 0 < y < 2, \\ \frac{dy}{dt} &< 0 && \text{when } -\infty < y < -2 \text{ or } 2 < y < \infty. \end{aligned}$$

The phase-line for this equation is therefore



- (b) If $y(0) = 1$, how does the solution $y(t)$ behave as $t \rightarrow \infty$?

Solution: It is clear from the answer to (a) that

$$\frac{dy}{dt} > 0 \quad \text{when } 0 < y < 2,$$

so that $y(t) \rightarrow 2$ as $t \rightarrow \infty$ if $y(0) = 1$.

- (c) If $y(0) = -1$, how does the solution $y(t)$ behave as $t \rightarrow \infty$?

Solution: It is clear from the answer to (a) that

$$\frac{dy}{dt} > 0 \quad \text{when } -2 < y < 0,$$

so that $y(t) \rightarrow 0$ as $t \rightarrow \infty$ if $y(0) = -1$.

- (d) Sketch a graph of y versus t showing the direction field and several solution curves. The graph should show all the stationary solutions as well as solution curves above and below each of them. Every value of y should lie on at least one sketched solution curve.

Solution: Will be given during the review session.

- (4) A tank initially contains 100 liters of pure water. Beginning at time $t = 0$ brine (salt water) with a salt concentration of 2 grams per liter (g/l) flows into the tank at a constant rate of 3 liters per minute (l/min) and the well-stirred mixture flows out of the tank at the same rate. Let $S(t)$ denote the mass (g) of salt in the tank at time $t \geq 0$.

- (a) Write down an initial-value problem that governs $S(t)$.

Solution: Because water flows in and out of the tank at the same rate, the tank will contain 100 liters of salt water for every $t > 0$. The salt concentration of the water in the tank at time t will therefore be $S(t)/100$ g/l. Because this is also the concentration of the outflow, $S(t)$, the mass of salt in the tank at time t , will satisfy

$$\frac{dS}{dt} = \text{RATE IN} - \text{RATE OUT} = 2 \cdot 3 - \frac{S}{100} \cdot 3 = 6 - \frac{3}{100}S.$$

Because there is no salt in the tank initially, the initial-value problem that governs $S(t)$ is

$$\frac{dS}{dt} = 6 - \frac{3}{100}S, \quad S(0) = 0.$$

- (b) Is $S(t)$ an increasing or decreasing function of t ? (Give your reasoning.)

Solution: One sees from part (a) that

$$\frac{dS}{dt} = \frac{3}{100}(200 - S) > 0 \quad \text{for } S < 200,$$

whereby $S(t)$ is an increasing function of t that will approach the stationary value of 200 g as $t \rightarrow \infty$.

- (c) What is the behavior of $S(t)$ as $t \rightarrow \infty$? (Give your reasoning.)

Solution: The argument given for part (b) already shows that $S(t)$ is an increasing function of t that approaches the stationary value of 200 g as $t \rightarrow \infty$.

(d) Derive an explicit formula for $S(t)$.

Solution: The differential equation given in the answer to part (a) is linear, so write it in the form

$$\frac{dS}{dt} + \frac{3}{100}S = 6.$$

An integrating factor is $e^{\frac{3}{100}t}$, whereby

$$\frac{d}{dt}(e^{\frac{3}{100}t}S) = 6e^{\frac{3}{100}t}.$$

This is then integrated to obtain

$$e^{\frac{3}{100}t}S = 200e^{\frac{3}{100}t} + C.$$

The integration constant C is found by setting $t = 0$ and $S = 0$, whereby

$$C = e^0 \cdot 0 - 200 \cdot e^0 = -200.$$

Then solving for S gives

$$S(t) = 200 - 200e^{-\frac{3}{100}t}.$$

(5) Suppose you are using the Heun-midpoint method to numerically approximate the solution of an initial-value problem over the time interval $[0, 5]$. By what factor would you expect the error to decrease when you increase the number of time steps taken from 500 to 2000.

Solution: The Heun-midpoint method is second order, which means its (global) error scales like h^2 where h is the time step. When the number of time steps taken increases from 500 to 2000, the time step h decreases by a factor of 4. The error will therefore decrease (like h^2) by a factor of $4^2 = 16$.

(6) Give an implicit general solution to each of the following differential equations.

(a) $\left(\frac{y}{x} + 3x\right) dx + (\log(x) - y) dy = 0.$

Solution: Because

$$\partial_y\left(\frac{y}{x} + 3x\right) = \frac{1}{x} = \partial_x(\log(x) - y) = \frac{1}{x},$$

the equation is *exact*. You can therefore find $H(x, y)$ such that

$$\partial_x H(x, y) = \frac{y}{x} + 3x, \quad \partial_y H(x, y) = \log(x) - y.$$

The first of these equations implies that

$$H(x, y) = y \log(x) + \frac{3}{2}x^2 + h(y).$$

Plugging this into the second equation then shows that

$$\log(x) - y = \partial_y H(x, y) = \log(x) + h'(y).$$

Hence, $h'(y) = -y$, which yields $h(y) = -\frac{1}{2}y^2$. The general solution is therefore governed implicitly by

$$y \log(x) + \frac{3}{2}x^2 - \frac{1}{2}y^2 = C,$$

where C is an arbitrary constant.

(b) $(x^2 + y^3 + 2x) dx + 3y^2 dy = 0$.

Solution: Because

$$\partial_y(x^2 + y^3 + 2x) = 3y^2 \neq \partial_x(3y^2) = 0,$$

the equation is *not exact*. Seek an integrating factor $\mu(x, y)$ such that

$$\partial_y((x^2 + y^3 + 2x)\mu) = \partial_x(3y^2\mu).$$

This means that μ must satisfy

$$(x^2 + y^3 + 2x)\partial_y\mu + 3y^2\mu = 3y^2\partial_x\mu.$$

If you assume that μ depends only on x (so that $\partial_y\mu = 0$) then this reduces to

$$\mu = \partial_x\mu,$$

which depends only on x . One sees from this that $\mu = e^x$ is an integrating factor. This implies that

$$(x^2 + y^3 + 2x)e^x dx + 3y^2e^x dy = 0 \quad \text{is exact.}$$

You can therefore find $H(x, y)$ such that

$$\partial_x H(x, y) = (x^2 + y^3 + 2x)e^x, \quad \partial_y H(x, y) = 3y^2e^x.$$

The second of these equations implies that

$$H(x, y) = y^3e^x + h(x).$$

Plugging this into the first equation then yields

$$(x^2 + y^3 + 2x)e^x = \partial_x H(x, y) = y^3e^x + h'(x).$$

Hence, h satisfies

$$h'(x) = (x^2 + 2x)e^x.$$

This can be integrated to obtain $h(x) = x^2e^x$. The general solution is therefore governed implicitly by

$$(y^3 + x^2)e^x = C,$$

where C is an arbitrary constant.

- (7) A 2 kilogram (kg) mass initially at rest is dropped in a medium that offers a resistance of $v^2/40$ newtons ($= \text{kg m/sec}^2$) where v is the downward velocity (m/sec) of the mass. The gravitational acceleration is 9.8 m/sec^2 .

(a) What is the terminal velocity of the mass?

Solution: The terminal velocity is the velocity at which the force of resistance balances that of gravity. This happens when

$$\frac{1}{40}v^2 = mg = 2 \cdot 9.8.$$

Upon solving this for v one obtains

$$\begin{aligned} v &= \sqrt{40 \cdot 2 \cdot 9.8} \text{ m/sec} && \text{(full marks)} \\ &= \sqrt{4 \cdot 2 \cdot 98} = \sqrt{4 \cdot 2 \cdot 2 \cdot 49} \\ &= \sqrt{4^2 \cdot 7^2} = 4 \cdot 7 = 28 \text{ m/sec.} \end{aligned}$$

- (b) Write down an initial-value problem that governs v as a function of time. (You do not have to solve it!)

Solution: The net downward force on the falling mass is the force of gravity minus the force of resistance. By Newton ($ma = F$), this leads to

$$m \frac{dv}{dt} = mg - \frac{1}{40}v^2.$$

Because $m = 2$ and $g = 9.8$, and because the mass is initially at rest, this yields the initial-value problem

$$\frac{dv}{dt} = 9.8 - \frac{1}{80}v^2, \quad v(0) = 0.$$

- (8) Consider the following MATLAB function M-file.

```
function [t,y] = solveit(ti, yi, tf, n)

h = (tf - ti)/n;
t = zeros(n + 1, 1);
y = zeros(n + 1, 1);
t(1) = ti;
y(1) = yi;
for i = 1:n
z = t(i)^4 + y(i)^2;
t(i + 1) = t(i) + h;
y(i + 1) = y(i) + (h/2)*(z + t(i + 1)^4 + (y(i) + h*z)^2);
end
```

- (a) What is the initial-value problem being approximated numerically?

Solution: The initial-value problem being approximated is

$$\frac{dy}{dt} = t^4 + y^2, \quad y(t_0) = y_0.$$

- (b) What is the numerical method being used?

Solution: The Heun-Trapezoidal (improved Euler) method is being used.

- (c) What are the output values of $t(2)$ and $y(2)$ that you would expect for input values of $t_i = 1$, $y_i = 1$, $t_f = 5$, $n = 20$?

Solution: The time step is given by $h = (t_f - t_i)/n = (5 - 1)/20 = 1/5 = .2$. The initial time and data are given by $t(1) = t_i = 1$ and $y(1) = y_i = 1$. One then has

$$\begin{aligned} t(2) &= t(1) + h = 1 + .2 = 1.2, \\ z &= t(1)^4 + y(1)^2 = 1 + 1 = 2, \\ y(2) &= y(1) + (h/2) (z + t(2)^4 + (y(1) + h z)^2) \\ &= 1 + .1(2 + (1.2)^4 + (1 + .2 \cdot 2)^2). \end{aligned}$$