

Advanced Calculus: MATH 410

Real Numbers

Professor David Levermore

7 January 2012

1. REAL NUMBER SYSTEM

1.1. Introduction. Numbers are at the heart of mathematics. By now you must be fairly familiar with them. Some basic sets of numbers are:

natural numbers, $\mathbb{N} = \{0, 1, 2, \dots\}$;

integers (die Zahlen), $\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$;

rational numbers (quotients), $\mathbb{Q} = \left\{ \frac{p}{q} : p, q \in \mathbb{Z}, q \neq 0 \right\}$;

real numbers, $\mathbb{R} = (-\infty, \infty)$;

complex numbers, $\mathbb{C} = \{x + iy : x, y \in \mathbb{R}\}$.

Each of these sets is endowed with natural algebraic operations (like ‘addition’ and ‘multiplication’) and order relations (like ‘less than’) by which their elements are manipulated and compared. It is fairly clear how \mathbb{N} , \mathbb{Z} , and \mathbb{Q} are related through an increasingly richer algebraic structure. It is also fairly clear that \mathbb{R} and \mathbb{C} bear a similar relationship. What is less clear is the relationship between \mathbb{Q} and \mathbb{R} . In particular, what are the properties that allow \mathbb{R} and not \mathbb{Q} to be identified with a ‘line’? In this section we address some of these issues.

We begin by addressing the question of why the rational numbers are inadequate for mathematical analysis. Simply put, the rationals do not allow us to solve equations that we would like to solve. This was also the reason behind the introduction of the negative integers and the rational numbers. The negative integers allow us to solve equations like $x + m = n$, where $m, n \in \mathbb{N}$. The rationals allow us to solve equations like $mx = n$, where $m, n \in \mathbb{Z}$ with $m \neq 0$. However, the rationals do not allow us to solve the rather simple equation $x^2 = 2$.

Proposition 1.1. *There exists no $x \in \mathbb{Q}$ such that $x^2 = 2$.*

Proof. We argue by assuming the contrary, and showing that it leads to a contradiction. Suppose there is such an $x \in \mathbb{Q}$. Then we can write it as $x = p/q$, where p and q are nonzero integers with no common factors. Because $x^2 = 2$, we see that $p^2 = 2q^2$. Hence, 2 is a factor of p^2 , which implies that 2 must also be a factor of p . We can therefore set $p = 2r$ for some nonzero integer r . Because $p^2 = 2q^2$, we see that $2^2r^2 = 2q^2$, which is the same as $q^2 = 2r^2$. Hence, 2 is a factor of q^2 , which implies that 2 must also be a factor of q . It follows that 2 is a factor of both p and q , which contradicts our assumption that p and q have no common factors. Therefore no such $x \in \mathbb{Q}$ exists. \square

There is nothing special about 2 in our argument. The same result is obtained for equations like $x^2 = n$ where n is any positive integer that is not a perfect square. More generally, the same result is obtained for equations like $x^m = n$ where m and n are positive integers such that $n \neq k^m$ for some integer k . The problem is that there are too many ‘holes’ like this in \mathbb{Q} . In this section we will see how \mathbb{R} fills these holes so as to allow the solution of such equations.

1.2. Fields. The sets \mathbb{Q} , \mathbb{R} , and \mathbb{C} endowed with their natural algebraic operations are each an example of a general algebraic structure known as a *field*.

Definition 1.1. A field is a set X equipped with two distinguished binary operations, called addition and multiplication, that satisfy the addition, multiplication, and distributive axioms presented below. Taken together, these axioms constitute the so-called field axioms.

Addition axioms. Addition maps any two $x, y \in X$ to their *sum* $x + y \in X$ such that:

- A1: $x + y = y + x \ \forall x, y \in X$, — commutativity;
- A2: $(x + y) + z = x + (y + z) \ \forall x, y, z \in X$, — associativity;
- A3: $\exists 0 \in X$ such that $x + 0 = x \ \forall x \in X$, — identity;
- A4: $\forall x \in X \ \exists -x \in X$, such that $x + (-x) = 0$, — inverse.

Multiplication axioms. Multiplication maps any two $x, y \in X$ to their *product* $xy \in X$ such that:

- M1: $xy = yx \ \forall x, y \in X$, — commutativity;
- M2: $(xy)z = x(yz) \ \forall x, y, z \in X$, — associativity;
- M3: $\exists 1 \in X$ such that $1 \neq 0$ and $x1 = x \ \forall x \in X$, — identity;
- M4: $\forall x \in X$ such that $x \neq 0 \ \exists x^{-1} \in X$ such that $xx^{-1} = 1$, — inverse.

Distributive axiom. Addition and multiplication are related by:

- D: $x(y + z) = xy + xz \ \forall x, y, z \in X$, — distributivity.

Examples. When addition and multiplication have their usual meaning, the field axioms clearly hold in \mathbb{Q} , \mathbb{R} , and \mathbb{C} , but not in \mathbb{N} or \mathbb{Z} . They also hold in $\mathbb{Z}_n \equiv \mathbb{Z}/(n\mathbb{Z})$ (the integers mod n) when n is prime. If you do not know this last example, do not worry. It is not critical in this course. You will see it in a basic algebra course.

All of the usual rules for algebraic manipulations involving addition, subtraction, multiplication, and division can be developed from the field axioms. This is not as easy as it sounds!

1.2.1. Consequences of the Addition Axioms. We begin by isolating the addition axioms.

Definition 1.2. A set X equipped with a distinguished binary operation that satisfies the addition axioms is called an Abelian group or a commutative group.

Examples. When addition has its usual meaning, the axioms for an Abelian groups clearly hold in \mathbb{Z} , \mathbb{Q} , \mathbb{R} , and \mathbb{C} , but not in \mathbb{N} . (As defined here, \mathbb{N} satisfies all these axioms but A4.) They also hold in \mathbb{Z}_n for every positive integer n .

The addition axioms immediately imply the following.

Proposition 1.2. Let X be an Abelian group.

- (a) If $x, y, z \in X$ and $x + y = x + z$ then $y = z$.
- (b) If $x, y \in X$ and $x + y = x$ then $y = 0$.
- (c) If $x, y \in X$ and $x + y = 0$ then $y = -x$.
- (d) If $x, y \in X$ then $-(x + y) = (-x) + (-y)$.
- (e) If $x \in X$ then $-(-x) = x$.

Proof. Exercise.

Assertion (a) states that addition enjoys a so-called cancellation law. Assertion (b) states that there is a unique additive identity of the type assumed in A3. This unique additive identity

is called *zero*. All other elements of X are said to be *nonzero*. Assertion (c) states that for every $x \in X$ there is a unique additive inverse of the type assumed in A4. This unique additive inverse is called the *negative* of x . The map defined for every $x \in X$ by $x \mapsto -x$ is called negation. Assertion (d) states that the negative of a sum is the sum of the negatives. Assertion (e) states that for every $x \in X$ the negative of the negative of x is again x .

When working with Abelian groups, it is both convenient and common to write

$$x - y, \quad x + y + z, \quad 2x, \quad 3x, \quad \dots,$$

rather than

$$x + (-y), \quad x + (y + z), \quad x + x, \quad x + x + x, \quad \dots.$$

More precisely, the symbol nx can be defined for every group element x and every natural number n by induction. We set $0x = 0$, where the second 0 is the additive identity, and define $(n + 1)x = nx + x$ for every $n \in \mathbb{N}$. This notation satisfies the following properties.

Proposition 1.3. *Let X be an Abelian group.*

- (a) *If $x \in X$ and $m, n \in \mathbb{N}$ then $(m + n)x = mx + nx$ and $(mn)x = n(mx)$.*
- (b) *If $x, y \in X$ and $n \in \mathbb{N}$ then $n(x + y) = nx + ny$.*
- (c) *If $x \in X$ and $n \in \mathbb{N}$ then $n(-x) = -(nx)$.*

Proof. Exercise.

Motivated by these facts, for every group element x the definition of the symbol nx can be extended to every integer n by setting $nx = (-n)(-x)$ when n is negative.

1.2.2. *Consequences of the Multiplication Axioms.* The only connection of the multiplication axioms to addition is through the references to zero in M3 and M4. An immediate consequence of M3 is that every field has at least two elements — 0 and 1. It is also clear that the nonzero elements of a field considered with the operation of multiplication form an Abelian group.

Examples. When addition and multiplication have their usual meaning, the addition and multiplication axioms clearly hold in \mathbb{Q} , \mathbb{R} , and \mathbb{C} , but not in \mathbb{N} or \mathbb{Z} . They also hold in \mathbb{Z}_n when n is prime.

The multiplication axioms immediately imply the following.

Proposition 1.4. *Let X be a field.*

- (a) *If $x, y, z \in X$, $x \neq 0$, and $xy = xz$ then $y = z$.*
- (b) *If $x, y \in X$, $x \neq 0$, and $xy = x$ then $y = 1$.*
- (c) *If $x, y \in X$, $x \neq 0$, and $xy = 1$ then $y = x^{-1}$.*
- (d) *If $x, y \in X$, $x \neq 0$ and $y \neq 0$ then $xy \neq 0$ and $(xy)^{-1} = x^{-1}y^{-1}$.*
- (e) *If $x \in X$ and $x \neq 0$ then $(x^{-1})^{-1} = x$.*

Proof. Exercise.

Assertion (a) states that multiplication enjoys a so-called cancellation law. Assertion (b) states that there is a unique multiplicative identity of the type assumed in M3. This unique multiplicative identity is called *one*. Assertion (c) states that for every nonzero $x \in X$ there is a unique multiplicative inverse of the type assumed in M4. This unique multiplicative inverse is called the *reciprocal* of x . The map defined for every nonzero $x \in X$ by $x \mapsto x^{-1}$ is called reciprocation. Assertion (d) states that the reciprocal of a product is the product of the reciprocals. Assertion (e) states that every nonzero $x \in X$ is the reciprocal of its reciprocal.

When working with fields, it is both convenient and common to write

$$x/y, \quad xyz, \quad x^2, \quad x^3, \quad \dots,$$

rather than

$$xy^{-1}, \quad x(yz), \quad xx, \quad xxx, \quad \dots.$$

More precisely, the symbol x^n can be defined for every field element x and every positive integer n by induction. We set $x^1 = x$ and define $x^{n+1} = x^n x$ for every $n \in \mathbb{Z}_+$, where \mathbb{Z}_+ denotes the positive integers. This notation satisfies the following properties.

Proposition 1.5. *Let X be a field.*

- (a) *If $x \in X$ and $m, n \in \mathbb{Z}_+$ then $x^{m+n} = x^m x^n$ and $x^{mn} = (x^m)^n$.*
- (b) *If $x, y \in X$ and $n \in \mathbb{Z}_+$ then $(xy)^n = x^n y^n$.*
- (c) *If $x \in X$, $x \neq 0$, and $n \in \mathbb{Z}_+$ then $x^n \neq 0$ and $(x^n)^{-1} = (x^{-1})^n$.*

Proof. Exercise.

Motivated by these facts, for every nonzero field element x the definition of the symbol x^n can be extended to every integer n by setting $x^0 = 1$, where the 1 is the multiplicative identity, and $x^n = (x^{-1})^{-n}$ when n is negative. The symbol 0^n remains undefined when n is not positive.

Exercise. Let X be a field. Extend Proposition 1.5 to \mathbb{Z} by proving the following.

- (a) If $x \in X$, $x \neq 0$, and $m, n \in \mathbb{Z}$ then $x^{m+n} = x^m x^n$ and $x^{mn} = (x^m)^n$.
- (b) If $x, y \in X$, $x \neq 0$, $y \neq 0$, and $n \in \mathbb{Z}$ then $(xy)^n = x^n y^n$.
- (c) If $x \in X$, $x \neq 0$, and $n \in \mathbb{Z}$ then $x^n \neq 0$ and $(x^n)^{-1} = (x^{-1})^n$.

1.2.3. *Consequences of the Distributive Axiom.* The distributive axiom gives the key connection between addition and multiplication. Taken together, the field axioms imply the following.

Proposition 1.6. *Let X be a field.*

- (a) *If $x \in X$ then $x0 = 0$.*
- (b) *If $x, y \in X$ and $xy = 0$ then $x = 0$ or $y = 0$.*
- (c) *If $x, y \in X$ then $(-x)y = -(xy) = x(-y)$.*
- (d) *If $x \in X$ and $x \neq 0$ then $(-x)^{-1} = -x^{-1}$.*

Proof. Exercise.

Assertion (a) states that the product of anything with zero is zero. In particular, it shows that zero cannot have a multiplicative inverse. Hence, an element has a multiplicative inverse if and only if it is nonzero. Assertion (b) states that if a product is zero, at least one of its factors must be zero. This should be compared with (d) of Proposition 2.2. Assertions (c) and (d) state how negation, multiplication, and reciprocation relate.

The field axioms allow you to extend to any field many of the formulas that you have known for years in the context of \mathbb{R} or \mathbb{C} . For example, you can establish the following formulas.

Proposition 1.7. *Let X be a field. Then for every $x, y \in X$ and every $n \in \mathbb{N}$ we have the difference of powers and binomial formulas*

$$(1.1) \quad x^{n+1} - y^{n+1} = (x - y) (x^n + x^{n-1}y + \dots + x^{n-k}y^k + \dots + xy^{n-1} + y^n),$$

$$(1.2) \quad (x + y)^n = x^n + nx^{n-1}y + \dots + \frac{n!}{(n-k)!k!} x^{n-k}y^k + \dots + nxy^{n-1} + y^n.$$

Proof. Exercise.

1.3. Ordered Sets. The sets \mathbb{N} , \mathbb{Z} , \mathbb{Q} , and \mathbb{R} endowed with their natural order relation are each an example of a general structure known as an ordered set.

Definition 1.3. An ordered set $(X, <)$ is a set X equipped with a distinguished binary relation “ $<$ ”, called an order, that satisfies the order axioms presented below.

Order axioms. A binary relation “ $<$ ” on a set X is called an *order* whenever:

- O1: if $x, y, z \in X$ then $x < y$ and $y < z$ implies $x < z$, — transitivity;
 O2: if $x, y \in X$ then exactly one of $x < y$, $x = y$, or $y < x$ is true, — trichotomy.

Examples. When “ $<$ ” has its usual meaning of “less than”, the order axioms clearly hold in \mathbb{N} , \mathbb{Z} , \mathbb{Q} , and \mathbb{R} . When “ $<$ ” has the unusual meaning of “greater than”, the order axioms also clearly hold in \mathbb{N} , \mathbb{Z} , \mathbb{Q} , and \mathbb{R} . We will stick with the usual meaning of “ $<$ ” in what follows.

When working with ordered sets, it is both convenient and common to use the notation

$$\begin{array}{lll} x > y, & x \leq y, & x \geq y, \\ x < y < z, & x < y \leq z, & \cdots, \end{array}$$

to respectively mean

$$\begin{array}{lll} y < x, & x < y \text{ or } x = y, & y < x \text{ or } x = y, \\ x < y \text{ and } y < z, & x < y \text{ and } y \leq z, & \cdots. \end{array}$$

1.3.1. Bounds. Ordered sets have associated notions of boundedness.

Definition 1.4. Let $(X, <)$ be an ordered set. A point $x \in X$ is an upper bound (a lower bound) of a set $S \subset X$ whenever $y \leq x$ ($x \leq y$) for every $y \in S$. If $S \subset X$ has an upper bound (a lower bound) then S is said to be bounded above (bounded below). A set $S \subset X$ that is both bounded above and bounded below is said to be bounded.

Definition 1.5. Let $(X, <)$ be an ordered set, and let $S \subset X$ be bounded above. A point $x \in X$ is a least upper bound or supremum of S whenever:

- (i) x is an upper bound of S ;
 (ii) if $y \in X$ is also an upper bound of S then $x \leq y$.

We similarly define a greatest lower bound or infimum of S .

If a supremum or infimum of S exists then it must be unique. The supremum of S is denoted $\sup\{S\}$ or $\sup\{z : z \in S\}$, while the infimum is denoted $\inf\{S\}$ or $\inf\{z : z \in S\}$. These notions should not be confused with those of maximum and minimum.

Definition 1.6. Let $(X, <)$ be an ordered set, and let $S \subset X$. A point $x \in S$ is a maximum (minimum) of S whenever x is an upper (lower) bound of S .

If a maximum or minimum of S exists then it must be unique. The maximum of S is denoted $\max\{S\}$ or $\max\{z : z \in S\}$, while the minimum is denoted $\min\{S\}$ or $\min\{z : z \in S\}$. Moreover, if a maximum (minimum) of S exists then

$$\sup\{S\} = \max\{S\} \quad (\inf\{S\} = \min\{S\}).$$

Examples. Any bounded open interval (a, b) in \mathbb{R} has neither a maximum nor a minimum, yet $\sup\{(a, b)\} = b$ and $\inf\{(a, b)\} = a$. For any bounded closed interval $[a, b]$ in \mathbb{R} one has $\max\{[a, b]\} = b$ and $\min\{[a, b]\} = a$. The same is true if these intervals and their endpoints are restricted to elements of \mathbb{Q} .

1.3.2. *Least Upper Bound Property.* What distinguishes \mathbb{R} from \mathbb{Q} is the following property.

Definition 1.7. *Let $(X, <)$ be an ordered set. Then X is said to have the least upper bound property whenever every nonempty subset of X with an upper bound has a least upper bound.*

Remark. It may seem we should also define a “greatest lower bound property”, but the next proposition shows that this is unnecessary because it is exactly the same property.

Proposition 1.8. *Let $(X, <)$ be an ordered set. Let X have the least upper bound property. Then every nonempty subset of X with a lower bound has a greatest lower bound.*

Proof. Let $S \subset X$ be a nonempty set with a lower bound. Let $L \subset X$ be the set of all lower bounds of S . It is nonempty and bounded above by any element of S . Therefore $\sup\{L\}$ exists. It is easy to check that $\sup\{L\} = \inf\{S\}$. \square

Examples. When “ $<$ ” has its usual meaning of “less than”, the sets \mathbb{N} and \mathbb{Z} have the least upper bound property. However, as we will show in the next proposition, the set \mathbb{Q} does not.

Proposition 1.9. *The set \mathbb{Q} does not have the least upper bound property.*

Proof. Consider the sets

$$S = \{r \in \mathbb{Q} : r > 0, r^2 < 2\}, \quad \tilde{S} = \{r \in \mathbb{Q} : r > 0, r^2 > 2\}.$$

These sets are clearly nonempty because $1 \in S$ and $2 \in \tilde{S}$. One can show that every point in \tilde{S} is an upper bound for S . In order to show that S has no least upper bound, one first shows that there is no $r \in \mathbb{Q}$ such that $r^2 = 2$. It follows (by trichotomy) that if p is a least upper bound of S then either $p \in S$ or $p \in \tilde{S}$. We will show that neither can be the case. More specifically, we will show that if $p \in S$ then p is not an upper bound of S , and that if $p \in \tilde{S}$ then p is not a least upper bound of S .

Let $p \in S$. We will construct a $q \in S$ such that $p < q$, thereby showing that p is not an upper bound of S . There are many ways to construct such a q . We are seeking a rational approximation of $\sqrt{2}$ from below that is better than p . This can be done by taking one iteration of Newton’s method applied to $f(x) = 1 - 2/x^2 = 0$. Set

$$q = p - \frac{f(p)}{f'(p)} = p - \frac{p^2 - 2}{4} p = \frac{6 - p^2}{4} p.$$

Because $x \mapsto f(x)$ is increasing and concave over $x > 0$, a picture alone should convince you this is a suitable q . Indeed, it is clear from the above formula that $0 < p < q$. A skeptic only needs to check that $q^2 < 2$. We confirm this fact by the calculation

$$2 - q^2 = 2 - \frac{36 - 12p^2 + p^4}{16} p^2 = \frac{(2 - p^2)^2(8 - p^2)}{16} > 0.$$

Now let $p \in \tilde{S}$. We will construct a $q \in \tilde{S}$ such that $q < p$, thereby showing that p is not a least upper bound of S . Once again, there are many ways to construct such a q . This time we are seeking a rational approximation of $\sqrt{2}$ from above that is better than p . This can be done by taking one iteration of Newton’s method and applied to $f(x) = x^2 - 2 = 0$. Set

$$q = p - \frac{f(p)}{f'(p)} = p - \frac{p^2 - 2}{2p} = \frac{p^2 + 2}{2p}.$$

Because $x \mapsto f(x)$ is increasing and convex over $x > 0$, a picture alone should convince you this is a suitable p . Indeed, it is clear from the above formula that $q < p$ and that $q > 0$. A skeptic only needs to check that $q^2 > 2$. We confirm this fact by the calculation

$$q^2 - 2 = \frac{p^4 + 4p^2 + 4}{4p^2} - 2 = \frac{p^4 - 4p^2 + 4}{4p^2} = \left(\frac{p^2 - 2}{2p}\right)^2 > 0.$$

□

Remark. An alternative construction that can be used for both cases in the above proof is

$$q = p - \frac{p^2 - 2}{p + 2} = \frac{2p + 2}{p + 2}.$$

Then

$$q^2 - 2 = \frac{4p^2 + 8p + 4}{p^2 + 4p + 4} - 2 = 2 \frac{p^2 - 2}{(p + 2)^2}.$$

While this construction yields a slicker proof, the underlying geometric picture seems less clear.

1.4. Ordered Fields. The sets \mathbb{Q} and \mathbb{R} endowed with their natural algebraic operations and order relation are each an example of a general algebraic structure known as an ordered field.

Definition 1.8. A set X that is both a field and an ordered set is called an ordered field whenever

OF1: if $x, y, z \in X$ then $x < y$ implies $x + z < y + z$;

OF2: if $x, y \in X$ then $0 < x$ and $0 < y$ implies $0 < xy$.

If $x > 0$ ($x < 0$, $x \geq 0$, $x \leq 0$) then we say x is positive (negative, nonnegative, nonpositive). The set of all positive (negative) elements of X is denoted X_+ (X_-).

Examples. When addition, multiplication, and “ $<$ ” have their usual meanings, the sets \mathbb{Q} and \mathbb{R} are ordered fields. In an algebra course you can learn that many other ordered fields arise in Galois theory.

1.4.1. Consequences of the Ordered Field Axioms. The ordered field axioms allow you to extend to any ordered field many of the rules for working with inequalities that you have known for years in the context of \mathbb{R} . For example, the rule that multiplying both sides of an inequality by a positive (negative) quantity will preserve (reverse) the inequality. Some of these rules are given in the following proposition.

Proposition 1.10. Let X be an ordered field.

- (a) If $x > 0$ then $-x < 0$, and vice versa.
- (b) If $x > 0$ and $y < z$ then $y < x + z$ and $xy < xz$.
- (c) If $x < 0$ and $y < z$ then $x + y < z$ and $xy > xz$.
- (d) If $x \neq 0$ then $x^2 > 0$.
- (e) If $0 < x < y$ and $n \in \mathbb{Z}_+$ then $0 < x^n < y^n$ and $0 < y^{-n} < x^{-n}$.

Proof. Exercise.

The above proposition shows that X_+ satisfies the following.

P1: If $x, y \in X_+$ then $x + y \in X_+$ and $xy \in X_+$.

P2: For every $x \in X$ exactly one of $x \in X_+$, $-x \in X_+$, or $x = 0$ is true.

These so-called positivity properties alone characterize the order relation on the field X .

Proposition 1.11. Let X be a field. Let $X_+ \subset X$ satisfy the positivity properties P1 and P2. Define the binary relation $<$ on X by

$$(1.3) \quad x < y \quad \text{means} \quad y - x \in X_+.$$

Then $(X, <)$ is an ordered field.

Proof. Exercise.

Remark. Proposition 1.11 implies that we could have defined an ordered field as a field X that has a subset X_+ satisfying the positivity properties, P1 and P2. In that case the positivity properties become the positivity axioms and, upon defining the order on X by (1.3), the order axioms O1, O2, OF1, and OF2 become order properties. This is the approach taken in Fitzpatrick's book.

1.4.2. *Absolute Value Function.* There is a natural absolute value function on any ordered field.

Definition 1.9. Let X be an ordered field. The absolute value function on X is defined by

$$|x| = \begin{cases} x & \text{if } x > 0 \\ 0 & \text{if } x = 0 \\ -x & \text{if } x < 0 \end{cases}.$$

Some of its properties are given in the following proposition. They should all look very familiar to you. However, your goal now is to understand how they follow from Definition 1.9 and the ordered field axioms.

Proposition 1.12. Let X be an ordered field. Then for every $x, y \in X$

- (a) $|x| \geq 0$, — nonnegativity;
- (b) $|x| = 0$ if and only if $x = 0$, — definiteness;
- (c) $|x + y| \leq |x| + |y|$, — triangle inequality;
- (d) $|xy| = |x||y|$, — multiplicativity;
- (e) $||x| - |y|| \leq |x - y|$, — difference inequality.

Proof. Exercise.

With the absolute value function we can define the *distance* between points $x, y \in X$ by $d(x, y) = |x - y|$. This distance function satisfies the following proposition.

Proposition 1.13. Let X be an ordered field. Let $d(x, y) = |x - y|$ where $|\cdot|$ is given by Definition 1.9. Then for every $x, y, z \in X$

- (a) $d(x, y) \geq 0$, — nonnegativity;
- (b) $d(x, y) = 0$ if and only if $x = y$, — definiteness;
- (c) $d(x, y) = d(y, x)$, — symmetry;
- (d) $d(x, z) \leq d(x, y) + d(y, z)$, — triangle inequality.

Proof. Exercise.

We can also characterize bounded sets with the absolute value function.

Proposition 1.14. Let X be an ordered field. Then $S \subset X$ is bounded if and only if there exists an $m \in X_+$ such that

$$x \in S \quad \implies \quad |x| \leq m.$$

Proof. Exercise.

1.5. Real Numbers. We now state without proof the main theorem of this section.

Theorem 1.1. *There exists a unique (up to an isomorphism) ordered field with the least upper bound property that contains \mathbb{Q} (up to an isomorphism) as a subfield.*

Proof. Proofs of this theorem are quite long and technical. You can find a proof of all but the uniqueness in the book by W. Rudin, *Principles of Mathematical Analysis*, McGraw-Hill, New York, 1976, and another in the book by T. Tao, *Analysis I*, Hindustan Book Agency, 2006 (available through the American Mathematical Society). The proof in Rudin is based upon a construction due to Richard Dedekind in which the real numbers are built from subsets of the rationals now called Dedekind cuts. The proof in Tao is based upon a construction due to Georg Cantor in which the real numbers are built from equivalence classes of Cauchy sequences within the rationals. Both Dedekind and Cantor published their constructions in 1872. \square

Definition 1.10. *The real numbers are defined to be the unique ordered field with the least upper bound property whose existence is guaranteed by Theorem 1.1. This field is denoted \mathbb{R} .*

Remark. The least upper bound property that sets \mathbb{R} apart from \mathbb{Q} . As we will see, it is why \mathbb{R} can be identified with a line.

1.5.1. Intervals. Intervals are special subsets of \mathbb{R} that will play a leading role in our study. They are denoted with the so-called interval notation. The empty set \emptyset is considered to be an interval. For every $a \in \mathbb{R}$ we define $[a, a] = \{a\}$. For every $a, b \in \mathbb{R}$ such that $a < b$ we define

$$\begin{aligned} (a, b) &= \{x \in \mathbb{R} : a < x < b\}, & [a, b) &= \{x \in \mathbb{R} : a \leq x < b\}, \\ [a, b] &= \{x \in \mathbb{R} : a < x \leq b\}, & [a, b] &= \{x \in \mathbb{R} : a \leq x \leq b\}. \end{aligned}$$

We call a and b respectively the *left endpoint* and the *right endpoint* of these intervals. Unless it is stated explicitly otherwise, when we write (a, b) , $[a, b)$, or $(a, b]$ it is implied that $a < b$, while when we write $[a, b]$ it is implied that $a \leq b$.

Similarly, for every $a, b \in \mathbb{R}$ we define

$$\begin{aligned} (a, \infty) &= \{x \in \mathbb{R} : a < x\}, & (-\infty, b) &= \{x \in \mathbb{R} : x < b\}, \\ [a, \infty) &= \{x \in \mathbb{R} : a \leq x\}, & (-\infty, b] &= \{x \in \mathbb{R} : x \leq b\}. \end{aligned}$$

We call a the *left endpoint* of (a, ∞) and $[a, \infty)$ and b the *right endpoint* of $(-\infty, b)$ and $(-\infty, b]$. Finally, we define $(-\infty, \infty) = \mathbb{R}$. Here we have not defined the symbols ∞ and $-\infty$ outside of the context of an unbounded interval.

An endpoint of an interval is said to be *closed* if it is contained in the interval, and is said to be *open* otherwise. More specifically, the point a is a closed endpoint of the intervals $[a, b]$, $[a, b)$, and $[a, \infty)$, while it is an open endpoint of $(a, b]$, (a, b) , and (a, ∞) . The point b is a closed endpoint of the intervals $[a, b]$, $(a, b]$, and $(-\infty, b]$, while it is an open endpoint of (a, b) , (a, b) , and $(-\infty, b)$.

An interval is said to be *closed* if all its endpoints are closed and is said to be *open* if all its endpoints are open. More specifically, $[a, b]$, $[a, \infty)$, and $(-\infty, b]$ are closed intervals, while (a, b) , (a, ∞) , and $(-\infty, b)$ are open intervals. The intervals \emptyset and $(-\infty, \infty)$ are both closed and open because they have no endpoints. Intervals $[a, b)$ and $(a, b]$ are neither closed nor open.

Exercise. Prove that for every $a \in \mathbb{R}$ and $r \in \mathbb{R}_+$ we have

$$\{x \in \mathbb{R} : |x - a| < r\} = (a - r, a + r), \quad \{x \in \mathbb{R} : |x - a| \leq r\} = [a - r, a + r].$$

1.5.2. *Properties of the Real Numbers.* The following important properties relate the reals \mathbb{R} with the positive integers \mathbb{Z}_+ , the integers \mathbb{Z} , and the rationals \mathbb{Q} .

Proposition 1.15. *The following hold.*

- If $x, y \in \mathbb{R}$ and $x > 0$ then there exists $n \in \mathbb{Z}_+$ such that

$$nx > y.$$

- If $x \in \mathbb{R}$ then there exists a unique $m \in \mathbb{Z}$ such that

$$m \in (x - 1, x] \quad (\text{or equivalently } x \in [m, m + 1)).$$

- If $x, y \in \mathbb{R}$ and $x < y$ then there exists a $q \in \mathbb{Q}$ such that

$$x < q < y.$$

Remark. The first assertion above is called the *Archimedean property* of \mathbb{R} , the second is a statement about the uniform distribution of the integers, while the third asserts that \mathbb{Q} is *dense* in \mathbb{R} — i.e. that between any two reals lies a rational.

Proof. Suppose the first assertion is false. Then the set $S = \{nx : n \in \mathbb{N}\}$ is bounded above by y . By the least upper bound property S has a supremum. Let $z = \sup\{S\}$. Because $x > 0$ one has that $z - x < z$. Hence, $z - x$ is not an upper bound for S because $z = \sup\{S\}$. This implies there exists some $n \in \mathbb{N}$ such that $z - x < nx$. But then $z < (n + 1)x$, which contradicts the fact z is an upper bound of S . Therefore the first assertion holds.

To prove the second assertion, by the first assertion there exists $k, l \in \mathbb{Z}_+$ such that $-x < k$ and $x < l$. It follows that $-k < x < l$. Because

$$x \in (-k, l) \subset \bigcup_{m=-k}^{l-1} [m, m + 1),$$

there exists some $m \in \mathbb{Z}$ such that

$$-k \leq m < l \quad \text{and} \quad m \leq x < m + 1.$$

It then follows that $m \in (x - 1, x]$. To prove uniqueness, suppose that $m, n \in \mathbb{Z}$ and $m, n \in (x - 1, x]$. Without loss of generality we may suppose that $m \leq n$ — otherwise simply exchange m and n . Because $x - 1 < m \leq n \leq x$, we see that $n - m \in \mathbb{Z}$ satisfies

$$0 \leq n - m < x - (x - 1) = 1.$$

It follows that $n - m = 0$, thereby establishing the uniqueness in the second assertion.

To prove the third assertion, because $y - x > 0$, by the first assertion there exists $n \in \mathbb{Z}_+$ such that $n(y - x) > 1$. Then by the second assertion there exists a unique $m \in (nx, nx + 1]$. Combining these facts yields

$$nx < m \leq nx + 1 < nx + n(y - x) = ny.$$

Because n is positive, we conclude that

$$x < \frac{m}{n} < y.$$

Therefore the third assertion holds with $q = m/n$. □

1.5.3. *Rational Powers.* Recall that we showed that $y^2 = 2$ had no solution within \mathbb{Q} . One of the most important consequences of the fact \mathbb{R} has least upper bound property is the existence of solutions to such equations.

Proposition 1.16. *For every $x \in \mathbb{R}_+$ and every $n \in \mathbb{Z}_+$ there exists a unique $y \in \mathbb{R}_+$ such that $y^n = x$.*

Proof. The uniqueness of such a y is clear because if $y < z$ then $y^n < z^n$. So we only have to show such a y exists. Consider the sets

$$S = \{r \in \mathbb{R}_+ : r^n < x\}, \quad \tilde{S} = \{r \in \mathbb{R}_+ : r^n > x\}.$$

The set S is nonempty because $s = x/(1+x) < 1$ implies $s^n < s < x$, whereby $s \in S$. The set \tilde{S} is nonempty because $1+x \in \tilde{S}$. One can show that every point in \tilde{S} is an upper bound for S . Let $y = \sup\{S\}$. Then (by trichotomy) either $y \in S$, $y \in \tilde{S}$, or $y^n = x$. We will show that the first two cases cannot occur, which will thereby prove the theorem. More specifically, we will show that no point in S is an upper bound of S , and that no point in \tilde{S} is the least upper bound of S .

Let $p \in S$. We can construct a $q \in S$ such that $p < q$, thereby showing that p is not an upper bound of S . This can be done by taking one iteration of Newton's method applied to $f(r) = 1 - x/r^n = 0$. The details are left as an exercise.

Now suppose $p \in \tilde{S}$. We can construct a $q \in \tilde{S}$ such that $q < p$, thereby showing that p is not a least upper bound of S . This can be done by taking one iteration of Newton's method and applied to $f(r) = r^n - x = 0$. The details are left as an exercise. \square

We designate the number y asserted in Proposition 1.16 by $x^{\frac{1}{n}}$.

Proposition 1.17. *Let $x \in \mathbb{R}_+$. Then*

$$(x^{\frac{1}{n}})^m = (x^m)^{\frac{1}{n}} \quad \text{for every } m \in \mathbb{Z} \text{ and } n \in \mathbb{Z}_+.$$

Proof. Just check that $(x^{\frac{1}{n}})^m$ and $(x^m)^{\frac{1}{n}}$ each solve the equation $y^n = x^m$. Equality then follows by Proposition 1.16. \square

By setting $x^{\frac{m}{n}} = (x^{\frac{1}{n}})^m = (x^m)^{\frac{1}{n}}$, we can therefore define x^p for every $x \in \mathbb{R}_+$ and $p \in \mathbb{Q}$. One can then show the following.

Proposition 1.18. *Let $x, y \in \mathbb{R}_+$ and $p, q \in \mathbb{Q}$. Then*

- (a) $x^{p+q} = x^p x^q$,
- (b) $(xy)^p = x^p y^p$,
- (c) $x^{pq} = (x^p)^q$, and $(x^p)^{-1} = (x^{-1})^p$.

Proof. Exercise.

1.6. Extended Real Numbers. It is convenient to extend the real numbers \mathbb{R} by appending two new elements designated $-\infty$ and ∞ . This enlarged set is called the extended real numbers and is denoted by \mathbb{R}_{ex} .

1.6.1. *Order.* The order $<$ on \mathbb{R} is extended to \mathbb{R}_{ex} by defining

$$-\infty < x < \infty \quad \text{for every } x \in \mathbb{R}.$$

The ordered set $(\mathbb{R}_{\text{ex}}, <)$ has the property that ∞ ($-\infty$) is an upper (lower) bound for every subset of \mathbb{R}_{ex} . It also has the least upper bound property. Indeed, the supremum of any $S \subset \mathbb{R}_{\text{ex}}$ is given by

$$\sup\{S\} = \begin{cases} \infty & \text{if } S \cap \mathbb{R} \text{ has no upper bound in } \mathbb{R} \text{ or } \infty \in S, \\ -\infty & \text{if } S = \{-\infty\} \text{ or } S = \emptyset, \\ \sup\{S \cap \mathbb{R}\} & \text{otherwise,} \end{cases}$$

where \emptyset denotes the empty set. In particular, every $S \subset \mathbb{R}$ that has no upper bound in \mathbb{R} (and therefore no supremum in \mathbb{R}) has $\sup\{S\} = \infty$ in \mathbb{R}_{ex} . Similar statements hold for lower bounds and infimums.

1.6.2. *Algebraic Operations.* The operations of addition and multiplication on R cannot be extended so as to make \mathbb{R}_{ex} into a field. It is however natural to extend addition by defining for every $x \in \mathbb{R}$

$$x + \infty = \infty + x = \infty, \quad x - \infty = -\infty + x = -\infty,$$

and by defining

$$\infty + \infty = \infty, \quad -\infty - \infty = -\infty,$$

while leaving $\infty - \infty$ and $-\infty + \infty$ undefined. In particular, $-\infty$ and ∞ do not have additive inverses.

Similarly, it is natural to extend multiplication by defining for every nonzero $x \in \mathbb{R}$

$$x \infty = \infty x = \begin{cases} \infty & \text{if } x > 0, \\ -\infty & \text{if } x < 0, \end{cases} \quad x(-\infty) = (-\infty)x = \begin{cases} -\infty & \text{if } x > 0, \\ \infty & \text{if } x < 0, \end{cases}$$

and by defining

$$\infty \infty = (-\infty)(-\infty) = \infty, \quad \infty(-\infty) = (-\infty)\infty = -\infty,$$

while leaving 0∞ , $\infty 0$, $0(-\infty)$, and $(-\infty)0$ undefined. In particular, 0 , $-\infty$ and ∞ do not have multiplicative inverses.

1.6.3. *Intervals.* Interval notation extends naturally to \mathbb{R}_{ex} . Given any $a \in \mathbb{R}_{\text{ex}}$ we define $[a, a] = \{a\}$. Given any $a, b \in \mathbb{R}_{\text{ex}}$ such that $a < b$ we define the sets

$$(1.4) \quad \begin{aligned} (a, b) &= \{x \in \mathbb{R}_{\text{ex}} : a < x < b\}, & [a, b) &= \{x \in \mathbb{R}_{\text{ex}} : a \leq x < b\}, \\ (a, b] &= \{x \in \mathbb{R}_{\text{ex}} : a < x \leq b\}, & [a, b] &= \{x \in \mathbb{R}_{\text{ex}} : a \leq x \leq b\}. \end{aligned}$$

When these sets are contained within \mathbb{R} the notation coincides with the interval notation we introduced earlier. We therefore call these sets *intervals* too. The new intervals are the ones that contain either $-\infty$ or ∞ — namely, ones that have the form $[-\infty, b)$, $[-\infty, b]$, $(a, \infty]$, or $[a, \infty]$ for some $a, b \in \mathbb{R}_{\text{ex}}$. In particular, one has $\mathbb{R}_{\text{ex}} = [-\infty, \infty]$.

We will often use the following characterization of intervals.

Proposition 1.19. (Interval Characterization Theorem.) *A set $S \subset \mathbb{R}_{\text{ex}}$ is an interval if and only if it has the property that*

$$(1.5) \quad \forall x, y \in \mathbb{R} \quad x, y \in S \quad \text{and} \quad x < y \quad \implies \quad (x, y) \subset S.$$

Proof. (\implies) It is clear from (1.4) that if S is an interval then it has property (1.5). In particular, the empty set and every singleton set (a set with only a single point in it) have property (1.5).

(\impliedby) The empty set and every singleton set is an interval. So we only have to consider sets that contain at least two points.

Let $S \subset \mathbb{R}_{\text{ex}}$ contain at least two points and have property (1.5). Because \mathbb{R}_{ex} has the least upper bound property, while every subset of \mathbb{R}_{ex} is bounded, we can set $a = \inf\{S\}$ and $b = \sup\{S\}$. You should be able to argue that $a < b$ because S has at least two points in it.

First, we show that $(a, b) \subset S$. Let $z \in (a, b)$. We claim that there exists $x \in (a, z)$ and $y \in (z, b)$ such that $x, y \in S$. (Otherwise z is either a lower or upper bound for S , which contradicts either $a = \inf\{S\}$ or $a = \sup\{S\}$.) Hence, property (1.5) implies that $z \in (x, y) \subset S$. Therefore, $(a, b) \subset S$.

Next, we claim that if $x < a$ or $x > b$ then $x \notin S$ because that would contradict either $a = \inf\{S\}$ or $a = \sup\{S\}$. When this fact is combined with the fact that $(a, b) \subset S$ it follows that S is an interval with

$$S = \begin{cases} (a, b) & \text{if } a \notin S \text{ and } b \notin S, \\ [a, b) & \text{if } a \in S \text{ and } b \notin S, \\ (a, b] & \text{if } a \notin S \text{ and } b \in S, \\ [a, b] & \text{if } a \in S \text{ and } b \in S, \end{cases}$$

where the sets (a, b) , $[a, b)$, $(a, b]$, and $[a, b]$ are defined by (1.4). □

2. SEQUENCES OF REAL NUMBERS

2.1. Sequences and Subsequences. Sequences play a central role in analysis. We introduce them here in the context of an arbitrary set X before specializing to sets of real numbers.

Definition 2.1. A sequence in a set X is a map from \mathbb{N} into X , often denoted $\{x_k\}$ or $\{x_k\}_{k \in \mathbb{N}}$, where $k \mapsto x_k$ maps the index $k \in \mathbb{N}$ to the point $x_k \in X$.

Remark. Any countable ordered set may be used as the index set instead of \mathbb{N} .

When X is an ordered set, sequences that either preserve or reverse order are special.

Definition 2.2. Let $(X, <)$ be an ordered set. A sequence $\{x_k\}_{k \in \mathbb{N}}$ in X is called

- increasing whenever $x_l > x_k$ for every $k, l \in \mathbb{N}$ with $l > k$,
- nondecreasing whenever $x_l \geq x_k$ for every $k, l \in \mathbb{N}$ with $l > k$,
- decreasing whenever $x_l < x_k$ for every $k, l \in \mathbb{N}$ with $l > k$,
- nonincreasing whenever $x_l \leq x_k$ for every $k, l \in \mathbb{N}$ with $l > k$.

It is called *monotonic* whenever it is either nondecreasing or nonincreasing. It is called *strictly monotonic* whenever it is either increasing or decreasing.

The following characterizations are often used to establish the monotonicity of a sequence.

Proposition 2.1. Let $(X, <)$ be an ordered set and $\{x_k\}_{k \in \mathbb{N}}$ be a sequence in X . The following characterizations hold.

| | | |
|---|----------------|---|
| $\{x_k\}_{k \in \mathbb{N}}$ is increasing | if and only if | $x_{k+1} > x_k$ for every $k \in \mathbb{N}$; |
| $\{x_k\}_{k \in \mathbb{N}}$ is nondecreasing | if and only if | $x_{k+1} \geq x_k$ for every $k \in \mathbb{N}$; |
| $\{x_k\}_{k \in \mathbb{N}}$ is decreasing | if and only if | $x_{k+1} < x_k$ for every $k \in \mathbb{N}$; |
| $\{x_k\}_{k \in \mathbb{N}}$ is nonincreasing | if and only if | $x_{k+1} \leq x_k$ for every $k \in \mathbb{N}$. |

Proof. Exercise. □

It is convenient to introduce the concepts of *eventually* and *frequently*.

Definition 2.3. Let $\mathcal{A}(x)$ be any assertion about any $x \in X$. (For example, $\mathcal{A}(x)$ could be the assertion “ $x \in S$ ” for a given $S \subset X$.) Let $\{x_k\}$ be a sequence in X . Then one says:

“ $\mathcal{A}(x_k)$ eventually as $k \rightarrow \infty$ ” when $\exists m \in \mathbb{N}$ such that $\forall k \geq m$ $\mathcal{A}(x_k)$;
“ $\mathcal{A}(x_k)$ frequently as $k \rightarrow \infty$ ” when $\forall m \in \mathbb{N} \exists k \geq m$ such that $\mathcal{A}(x_k)$.

When there is no possible confusion as to the index set, one says simply “ $\mathcal{A}(x_k)$ eventually” or “ $\mathcal{A}(x_k)$ frequently”, dropping the “as $k \rightarrow \infty$ ”.

Exercise. Show that $2^{-k} < .001$ eventually.

Exercise. Let $\{x_k\}$ be a sequence in X . Let $\mathcal{A}(x)$ be any assertion about any $x \in X$ and let $\sim \mathcal{A}(x)$ be its negation. Show that the negation of “ $\mathcal{A}(x_k)$ eventually” is “ $\sim \mathcal{A}(x_k)$ frequently”.

Exercise. Show that $\cos(k) > .5$ frequently, but not eventually.

Another useful concept is that of a *subsequence*.

Definition 2.4. A subsequence $\{x_{n_k}\}_{k \in \mathbb{N}}$ of a sequence $\{x_k\}_{k \in \mathbb{N}}$ in a set X is a map from \mathbb{N} into X given by $k \mapsto x_{n_k}$, where $\{n_k\}_{k \in \mathbb{N}}$ is an increasing sequence in \mathbb{N} .

Example. If $\{x_k\}$ is a sequence in a set X , then $\{x_{2k}\}$ is the subsequence with indices that are even, while $\{x_{k^3}\}$ is the subsequence with indices that are cubes.

Exercise. Consider the sequence $\{2^k\}$. Write out the first three terms (i.e. $k = 0, 1, 2$) in the subsequences $\{2^{3k}\}$ and $\{2^{2k+1}\}$.

In an ordered set, subsequences of monotonic sequences are again monotonic.

Proposition 2.2. *Let $(X, <)$ be an ordered set. Let $\{x_k\}_{k \in \mathbb{N}}$ be a sequence in X that is increasing (nondecreasing, decreasing, nonincreasing). Then every subsequence of $\{x_k\}_{k \in \mathbb{N}}$ is also increasing (nondecreasing, decreasing, nonincreasing).*

Proof. Exercise.

You should test your understanding of the concepts in this section by proving the following.

Proposition 2.3. *Let X be a set. Let $\mathcal{A}(x)$ be any assertion about any $x \in X$. Let $\{x_k\}_{k \in \mathbb{N}}$ be a sequence in X . Then $\mathcal{A}(x_k)$ frequently as $k \rightarrow \infty$ if and only if there exists a subsequence $\{x_{n_k}\}_{k \in \mathbb{N}}$ such that $\mathcal{A}(x_{n_k})$ eventually as $k \rightarrow \infty$.*

Proof. Exercise.

2.2. Convergence and Divergence. The most important concept related to sequences is that of convergence. Here we see it in the context of real sequences.

Definition 2.5. *A sequence $\{a_k\}_{k \in \mathbb{N}} \subset \mathbb{R}$ is said to converge or is said to be convergent whenever there exists a point $a \in \mathbb{R}$ such that for every $\epsilon > 0$ one has that*

$$(2.1) \quad |a_k - a| < \epsilon \text{ eventually as } k \rightarrow \infty.$$

This is denoted as

$$a_k \rightarrow a \text{ as } k \rightarrow \infty,$$

or as

$$\lim_{k \rightarrow \infty} a_k = a.$$

One then says that the sequence converges to a . A sequence that does not converge is said to diverge or is said to be divergent.

An immediate consequence of this definition and ideas from the previous section is the following proposition.

Proposition 2.4. *A sequence $\{a_k\}_{k \in \mathbb{N}} \subset \mathbb{R}$ diverges if and only if for every $a \in \mathbb{R}$ there exists an $\epsilon_a > 0$ such that*

$$|a_k - a| \geq \epsilon_a \text{ frequently as } k \rightarrow \infty.$$

Proof. Exercise.

Definition 2.5 does not assert that there is a unique number a that satisfies (2.1). The following proposition establishes this and more.

Proposition 2.5. *If a sequence $\{a_k\}_{k \in \mathbb{N}} \subset \mathbb{R}$ converges, there is a unique point in \mathbb{R} to which it converges. Moreover, the set $\{a_k\} \subset \mathbb{R}$ is bounded.*

Proof. Here we prove only the boundedness assertion. The proof of the uniqueness assertion is left as an exercise.

Let $\{a_k\}_{k \in \mathbb{N}}$ converge to $a \in \mathbb{R}$. Then $\exists m \in \mathbb{N}$ such that $\forall k \geq m$ $|a_k - a| < 1$. In particular, for every $k \geq m$ one has that $a - 1 < a_k < a + 1$. Then for every $k \in \mathbb{N}$ we have

$$|a_k| < 1 + \max\{|a_0|, |a_1|, \dots, |a_{m-1}|, |a|\}.$$

The sequence $\{a_k\}_{k \in \mathbb{N}}$ is therefore bounded. \square

Definition 2.6. *The unique point to which a convergent sequence in \mathbb{R} converges is called the limit of the sequence.*

An important characterization of the limit of a convergent sequence is given by the following.

Proposition 2.6. *Let $\{b_k\}_{k \in \mathbb{N}}$ be a convergent sequence in \mathbb{R} . Let $b \in \mathbb{R}$. Then*

$$\lim_{k \rightarrow \infty} b_k = b,$$

if and only if

$$a < b < c \quad \Longrightarrow \quad a < b_k < c \quad \text{eventually.}$$

Proof. Exercise.

Subsequences of a convergent sequence are also convergent, and have the same limit.

Proposition 2.7. *If a sequence $\{a_k\}_{k \in \mathbb{N}} \subset \mathbb{R}$ converges to a limit $a \in \mathbb{R}$ then every subsequence of $\{a_k\}_{k \in \mathbb{N}}$ also converges to a .*

Proof. Exercise.

The main theorem regarding basic operations, order, and limits is the following.

Proposition 2.8. *Let $\{a_k\}$ and $\{b_k\}$ be convergent sequences in \mathbb{R} with $a_k \rightarrow a$ and $b_k \rightarrow b$ as $k \rightarrow \infty$. Then*

- (i) $(a_k + b_k) \rightarrow (a + b)$,
- (ii) $-a_k \rightarrow -a$,
- (iii) $a_k b_k \rightarrow ab$,
- (iv) $1/a_k \rightarrow 1/a$ provided no division by zero occurs,
- (v) $|a_k| \rightarrow |a|$.

Moreover, if $a_k \leq b_k$ frequently then $a \leq b$. (Equivalently, if $a < b$ then $a_k < b_k$ eventually.)

Proof. Exercise.

When working with real sequences, it is useful to distinguish two of the many ways in which a sequence might diverge — namely, those when the sequence “approaches” either ∞ or $-\infty$.

Definition 2.7. *A sequence $\{a_k\}_{k \in \mathbb{N}} \subset \mathbb{R}$ is said to diverge to ∞ (to $-\infty$) if for every $b \in \mathbb{R}$ one has that*

$$a_k > b \text{ eventually (} a_k < b \text{ eventually) as } k \rightarrow \infty.$$

This is denoted as

$$a_k \rightarrow \infty \quad (a_k \rightarrow -\infty) \quad \text{as } k \rightarrow \infty,$$

or as

$$\lim_{k \rightarrow \infty} a_k = \infty \quad \left(\lim_{k \rightarrow \infty} a_k = -\infty \right).$$

One then says that the sequence approaches ∞ (approaches $-\infty$).

2.3. Monotonic Sequences. For monotonic sequences the least upper bound property can be employed to show the existence of limits.

Proposition 2.9. (Monotonic Sequence Theorem) *Let $\{a_k\}_{k \in \mathbb{N}}$ be a sequence in \mathbb{R} that is nondecreasing (nonincreasing). Then it converges if and only if it is bounded above (bounded below). Moreover, if it converges then*

$$\lim_{k \rightarrow \infty} a_k = \sup\{a_k\} \quad \left(\lim_{k \rightarrow \infty} a_k = \inf\{a_k\} \right),$$

while if it diverges then

$$\lim_{k \rightarrow \infty} a_k = \infty \quad \left(\lim_{k \rightarrow \infty} a_k = -\infty \right).$$

Proof. We give the proof for the nondecreasing case only; the nonincreasing case goes similarly. The proof of the assertion in the case of divergence in is left as an exercise.

(\Rightarrow) This follows from Proposition 2.5, which states that every convergent sequence is bounded.

(\Leftarrow) Because $\{a_k\}$ is bounded above, we can set $a = \sup\{a_k; k \in \mathbb{N}\}$ by the least upper bound property. We claim that $a_k \rightarrow a$ as $k \rightarrow \infty$. Let $\epsilon > 0$ be arbitrary. There exists some $m_\epsilon \in \mathbb{N}$ such that $0 \leq a - a_{m_\epsilon} < \epsilon$. (For if not, it would mean that $a_k \leq a - \epsilon$ for every $k \in \mathbb{N}$, which would contradict the definition of a .) But then for every $k > m_\epsilon$ one has that $0 \leq a - a_k \leq a - a_{m_\epsilon} < \epsilon$, which establishes the claim. \square

For monotonic sequences it is enough to know what happens to a single subsequence.

Proposition 2.10. *Let $\{a_k\}$ be a monotonic sequence in \mathbb{R} . Then $\{a_k\}$ is convergent if and only if it has a convergent subsequence.*

Proof. Exercise.

The Monotonic Sequence Theorem has the following consequence, often attributed to Cantor.

Proposition 2.11. (Nested Interval Theorem.) *Let $\{[a_k, b_k]\}_{k \in \mathbb{N}}$ be a sequence of closed, bounded intervals in \mathbb{R} that is nested in the sense that*

$$[a_{k+1}, b_{k+1}] \subset [a_k, b_k] \quad \text{for every } k \in \mathbb{N}.$$

Then the sequences $\{a_k\}$ and $\{b_k\}$ converge and

$$\bigcap_{k \in \mathbb{N}} [a_k, b_k] = [a, b], \quad \text{where } a = \lim_{k \rightarrow \infty} a_k, \quad b = \lim_{k \rightarrow \infty} b_k, \quad \text{with } a \leq b.$$

Proof. Because the sequence of intervals $\{[a_k, b_k]\}$ is nested, it follows that the sequence $\{a_k\}$ is nondecreasing while the sequence $\{b_k\}$ is nonincreasing. Hence, for every $m, n \in \mathbb{N}$ we have $a_m \leq a_n \leq b_n$ when $m \leq n$ and $a_m \leq b_m \leq b_n$ when $m \geq n$. This implies the sequence $\{a_k\}$ is bounded above by every b_n , while the sequence $\{b_k\}$ is bounded below by every a_n . The Monotone Sequence Theorem then implies that the sequences $\{a_k\}$ and $\{b_k\}$ converge with

$$a = \lim_{k \rightarrow \infty} a_k = \sup_k \{a_k\} \leq b_n, \quad b = \lim_{k \rightarrow \infty} b_k = \inf_k \{b_k\} \geq a_n.$$

Because $a_k \leq b_k$ for every $k \in \mathbb{N}$ it follows that $a \leq b$.

If $x \in [a, b]$ then because $[a, b] \subset [a_k, b_k]$ for every $k \in \mathbb{N}$, it follows that $x \in \bigcap_k [a_k, b_k]$. If $x < a$ then because $a_k \rightarrow a$, it follows that $x < a_k$ eventually, whereby $x \notin \bigcap_k [a_k, b_k]$. Similarly, if $b < x$ then because $b_k \rightarrow b$, it follows that $b_k < x$ eventually, whereby $x \notin \bigcap_k [a_k, b_k]$. \square

Remark. This theorem shows the above intersection of nested intervals is always nonempty. In particular, when $a = b$ this intersection consists of a single point. To better appreciate significance of this result, you should do the following exercise.

Exercise. Show that a nested sequence of closed, bounded intervals in \mathbb{Q} can have an empty intersection.

2.4. Limits and e . While Isaac Newton discovered e , it was Leonhard Euler who named it. Euler also showed that

$$\lim_{k \rightarrow \infty} \left(1 + \frac{1}{k}\right)^k = \lim_{k \rightarrow \infty} \left(1 - \frac{1}{k}\right)^{-k} = e.$$

Here we will use the Monotonic Sequence Theorem to prove that these limits exist. We will then consider e to be defined by these limits. In fact, we prove that more general limits exist.

Proposition 2.12. *For every $x \in \mathbb{R}$ we have the convergent limits*

$$(2.2) \quad \lim_{k \rightarrow \infty} \left(1 + \frac{x}{k}\right)^k = \lim_{k \rightarrow \infty} \left(1 - \frac{x}{k}\right)^{-k}.$$

Moreover, when $x \neq 0$ the sequence in the first limit is increasing for $k > -x$ while that in the second limit is decreasing for $k > x$.

Remark. Later we will see these limits converge to e^x . Here we only show they converge.

Proof. Let $x \in \mathbb{R}$. When $x = 0$ the result is easy because both sequences reduce to the constant sequence $\{1\}$, so we only need to consider the case $x \neq 0$.

We will use the following inequalities, which can be derived from the difference of powers formula (1.1). Specifically, for every $n \in \mathbb{Z}_+$ and every $z > y > 0$ we have the inequalities

$$(2.3) \quad (n+1)(z-y)y^n < z^{n+1} - y^{n+1} < (n+1)(z-y)z^n.$$

If $x > 0$ and $k \in \mathbb{Z}_+$ then setting $z = 1 + \frac{x}{k}$, $y = 1 + \frac{x}{k+1}$, and $n = k$ into the second inequality of (2.3) yields

$$\left(1 + \frac{x}{k}\right)^{k+1} - \left(1 + \frac{x}{k+1}\right)^{k+1} < \frac{x}{k} \left(1 + \frac{x}{k}\right)^k.$$

If $x < 0$ and $k \in \mathbb{Z}_+$ with $k > -x$ then setting $z = 1 + \frac{x}{k+1}$, $y = 1 + \frac{x}{k}$, and $n = k$ into the first inequality of (2.3) yields

$$-\frac{x}{k} \left(1 + \frac{x}{k}\right)^k < \left(1 + \frac{x}{k+1}\right)^{k+1} - \left(1 + \frac{x}{k}\right)^{k+1}.$$

By combining these two inequalities we see that

$$a_k = \left(1 + \frac{x}{k}\right)^k < \left(1 + \frac{x}{k+1}\right)^{k+1} = a_{k+1} \quad \text{for every } k > -x.$$

Upon replacing x with $-x$ and taking reciprocals we obtain

$$b_{k+1} = \left(1 - \frac{x}{k+1}\right)^{-(k+1)} < \left(1 - \frac{x}{k}\right)^{-k} = b_k \quad \text{for every } k > x.$$

Therefore $\{a_k\}$ is increasing for $k > -x$ while $\{b_k\}$ is decreasing for $k > x$.

Moreover, for $k > |x|$ we have

$$\frac{a_k}{b_k} = \left(1 + \frac{x}{k}\right)^k \left(1 - \frac{x}{k}\right)^k = \left(1 - \frac{x^2}{k^2}\right)^k < 1.$$

When this fact is combined with the monotonicity facts we conclude that

$$a_k < b_l \quad \text{for all integers } k, l \geq |x|.$$

Hence, $\{a_k\}$ is bounded above by every b_l while $\{b_l\}$ is bounded below by every a_k . By the Monotonic Sequence Theorem (Proposition 2.9) these sequences are convergent with

$$\lim_{k \rightarrow \infty} a_k \leq \lim_{l \rightarrow \infty} b_l.$$

If $k > 1$ then setting $z = 1$, $y = 1 - \frac{x^2}{k^2}$, and $n = k - 1$ into the second inequality of (2.3) yields

$$1 - \left(1 - \frac{x^2}{k^2}\right)^k < \frac{x^2}{k}.$$

Therefore for every $k > 1 + |x|$ we have

$$\left(1 - \frac{x^2}{k}\right) b_k < a_k < b_k.$$

By passing to the limit in these inequalities we see that $\{a_k\}$ and $\{b_k\}$ have the same limit. \square

2.5. Wallis Product Formula*. Soon after Newton developed calculus, his friend John Wallis showed that π could be expressed with the following limit.

Proposition 2.13.

$$(2.4) \quad \lim_{k \rightarrow \infty} \frac{2^2 \cdot 4^2 \cdot 6^2 \cdots (2k-2)^2 \cdot (2k)^2}{1 \cdot 3^2 \cdot 5^2 \cdots (2k-1)^2 \cdot (2k+1)} = \frac{\pi}{2}.$$

Proof. We will prove this limit after we borrow some facts from calculus that we have not covered yet in this course, but that you already know. For every $n \in \mathbb{N}$ we define

$$c_n = \int_0^{\frac{\pi}{2}} \cos(x)^n dx.$$

Because $0 \leq \cos(x) \leq 1$ for every $x \in [0, \frac{\pi}{2}]$ we see that $\{c_n\}_{n \in \mathbb{N}}$ is a nonincreasing sequence. We obtain $c_0 = \frac{\pi}{2}$ and $c_1 = 1$ by direct integration. We can use integration by parts to derive the recursion formula

$$c_n = \frac{n-1}{n} c_{n-2} \quad \text{for every } n \geq 2.$$

With these facts in hand, we can prove the Wallis product formula (2.4).

First, we can use induction to show that for every $k \in \mathbb{N}$

$$c_{2k} = \frac{\pi}{2} \cdot \frac{1}{2} \cdot \frac{3}{4} \cdot \frac{5}{6} \cdots \frac{2k-3}{2k-2} \cdot \frac{2k-1}{2k}, \quad c_{2k+1} = 1 \cdot \frac{2}{3} \cdot \frac{4}{5} \cdot \frac{6}{7} \cdots \frac{2k-2}{2k-1} \cdot \frac{2k}{2k+1}.$$

Because $\{c_n\}_{n \in \mathbb{N}}$ is a nonincreasing sequence, we have $c_{2k+2} \leq c_{2k+1} \leq c_{2k}$ for every $k \in \mathbb{N}$. Upon dividing this by c_{2k} and multiplying the result by $\frac{\pi}{2}$, we obtain the inequalities

$$\frac{2k+1}{2k+2} \cdot \frac{\pi}{2} \leq \frac{2^2 \cdot 4^2 \cdot 6^2 \cdots (2k-2)^2 \cdot (2k)^2}{1 \cdot 3^2 \cdot 5^2 \cdots (2k-1)^2 \cdot (2k+1)} \leq \frac{\pi}{2}.$$

By passing to the limit in these inequalities we obtain the Wallis product formula (2.4). \square

2.6. **De Moivre-Stirling Formula***. Another famous limit is the de Moivre-Stirling formula.

Proposition 2.14. (De Moivre-Stirling)

$$(2.5) \quad \lim_{n \rightarrow \infty} \frac{n!}{\sqrt{2\pi n} n^n e^{-n}} = 1.$$

Remark. This shows that $n!$ behaves like $\sqrt{2\pi n} (n/e)^n$ for large n . The convergence of the limit was established by Abarham de Moivre in his work on probability. John Stirling subsequently used the Wallis product formula to evaluate the limit. Formula (2.5) is commonly called the Stirling Formula or the Stirling Approximation, which are names that give Stirling all the credit for a result that was due largely to de Moivre.

We first use the Monotonic Sequence Theorem to show that the limit exists.

Proposition 2.15. (De Moivre Convergence Theorem)

$$(2.6) \quad \lim_{n \rightarrow \infty} \frac{n!}{\sqrt{n} n^n e^{-n}} \text{ converges to a positive limit.}$$

Moreover, the sequence that appears in the limit is decreasing.

Remark. The following proof uses facts from calculus that we have not covered yet. For example, it uses the functions $\log(x)$, e^x , and their basic properties. Strictly speaking, our proof will be incomplete until all these logical gaps are filled. However, we give it now to illustrate a substantial application of the Monotonic Sequence Theorem.

Proof. We will bound $\ell_n = \log(n!)$. By properties of logarithms

$$(2.7) \quad \ell_n = \log(n!) = \log(1 \cdot 2 \cdots (n-1) \cdot n) = \sum_{k=1}^n \log(k).$$

This looks like a numerical approximation to $\int_1^n \log(x) dx$. We will use the fact that $\log(x)$ is strictly concave (down) over $(0, \infty)$ and the fact that the trapezoidal approximation $\text{Trap}(h)$ with uniformly subintervals of length h underestimates the definite integral $\int_a^b f(x) dx$ of a differentiable strictly concave function $f(x)$ with a correction that satisfies the bounds

$$0 < \int_a^b f(x) dx - \text{Trap}(h) < \frac{h^2}{8} (f'(a) - f'(b)).$$

In particular, for $a = 1$, $b = n$, $f(x) = \log(x)$, and $h = 1$ we have $\text{Trap}(1) = \ell_n - \frac{1}{2} \log(n)$ and $f'(x) = 1/x$, whereby

$$(2.8) \quad 0 < \int_1^n \log(x) dx + \frac{1}{2} \log(n) - \ell_n < \frac{1}{8} \left(1 - \frac{1}{n}\right) < \frac{1}{8}.$$

Now define the sequence $\{c_n\}_{n=2}^\infty$ by

$$(2.9) \quad c_n = \int_1^n \log(x) dx + \frac{1}{2} \log(n) - \ell_n = \sum_{k=1}^{n-1} \left(\int_k^{k+1} \log(x) dx - \frac{\log(k) + \log(k+1)}{2} \right).$$

The terms in this sum are the corrections of the trapezoidal approximation to $\int_k^{k+1} \log(x) dx$ with $h = 1$, so they are each positive. The sequence $\{c_n\}_{n=2}^\infty$ is therefore increasing. It also is bounded with $0 < c_n < \frac{1}{8}$ by (2.8).

Because $x \log(x) - x$ is a primitive (antiderivative) of $\log(x)$, we have

$$c_n = \int_1^n \log(x) dx + \frac{1}{2} \log(n) - \ell_n = (n + \frac{1}{2}) \log(n) - n + 1 - \ell_n.$$

Therefore, by properties of exponentials and logarithms we have

$$(2.10) \quad r_n = \frac{n!}{\sqrt{n} n^n e^{-n}} = \exp(\ell_n - (n + \frac{1}{2}) \log(n) + n) = e^{1-c_n}.$$

Because $\{c_n\}_{n=2}^\infty$ is an increasing sequence that satisfies the bounds $0 < c_n < \frac{1}{8}$ while e^x is an increasing function of x , we see that $\{r_n\}_{n=2}^\infty$ is a decreasing sequence that satisfies the bounds

$$(2.11) \quad e^{\frac{7}{8}} < r_n = \frac{n!}{\sqrt{n} n^n e^{-n}} < e.$$

Therefore $\{r_n\}_{n=2}^\infty$ converges by the Monotonic Sequence Theorem (Proposition 2.9). Moreover, its limit lies within $[e^{\frac{7}{8}}, e]$ and is thereby positive. \square

We now prove Proposition 2.14 by evaluating the limit using the Wallis product formula.

Proof. Let r_n be defined by (2.10). By (2.11) and the de Moivre Convergence Theorem

$$\lim_{n \rightarrow \infty} r_n = r \quad \text{for some } r \in [e^{\frac{7}{8}}, e].$$

To establish (2.5) we need to show that $r = \sqrt{2\pi}$. The Wallis product formula (2.4) states

$$\lim_{n \rightarrow \infty} w_n = \frac{\pi}{2} \quad \text{where} \quad w_n = \frac{2^2 \cdot 4^2 \cdot 6^2 \cdots (2n-2)^2 \cdot (2n)^2}{1 \cdot 3^2 \cdot 5^2 \cdots (2n-1)^2 \cdot (2n+1)}.$$

It is easily checked that

$$\begin{aligned} \frac{r_n^2}{r_{2n}} &= \frac{(n!)^2}{n n^{2n} e^{-2n}} \cdot \frac{\sqrt{2n} (2n)^{2n} e^{-2n}}{(2n)!} = \sqrt{\frac{2}{n}} \cdot \frac{(2^n n!)^2}{(2n)!} \\ &= \sqrt{\frac{2}{n}} \cdot \frac{2 \cdot 4 \cdot 6 \cdots (2n-2) \cdot 2n}{1 \cdot 3 \cdot 5 \cdots (2n-3) \cdot (2n-1)} = \sqrt{\frac{2}{n}} (2n+1) w_n. \end{aligned}$$

Hence,

$$r = \lim_{n \rightarrow \infty} \frac{r_n^2}{r_{2n}} = \lim_{n \rightarrow \infty} \sqrt{\frac{2}{n}} (2n+1) w_n = \sqrt{4 \frac{\pi}{2}} = \sqrt{2\pi}.$$

Therefore the de Moivre-Stirling formula (2.5) holds. \square

Exercise. Prove the factorial-root limit

$$(2.12) \quad \lim_{n \rightarrow \infty} \frac{(n!)^{\frac{1}{n}}}{n} = \frac{1}{e}.$$

Exercise. Define the sequence $\{d_n\}_{n=2}^\infty$ by

$$d_n = \int_1^{n+\frac{1}{2}} \log(x) dx - \ell_n,$$

where ℓ_n is given by (2.7). Show that $\{d_n\}_{n=2}^\infty$ is a decreasing sequence such that $c_n < d_n$ for every $n \geq 2$, where c_n is given by (2.9). Show moreover that $\{d_n\}_{n=2}^\infty$ is convergent with

$$\lim_{n \rightarrow \infty} d_n = \lim_{n \rightarrow \infty} c_n.$$

2.7. Limit Superior and Limit Inferior. The power of the Monotonic Sequence Theorem (Proposition 2.9) lies in the fact that from every real sequence $\{a_k\}_{k \in \mathbb{N}}$ that is bounded above (bounded below), we can construct a nonincreasing (nondecreasing) sequence from its “tails”. Specifically, we construct the sequence $\{\bar{a}_k\}$ ($\{\underline{a}_k\}$) with elements defined by

$$\bar{a}_k = \sup\{a_l : l \geq k\} \quad (\underline{a}_k = \inf\{a_l : l \geq k\}) .$$

(Here you should think of \bar{a} and \underline{a} as new letters that are related to a .) This sequence is clearly nonincreasing (nondecreasing). The convergence of such sequences is characterized by the Monotonic Sequence Theorem, which motivates the following definition.

Definition 2.8. For every sequence $\{a_k\}_{k \in \mathbb{N}}$ in \mathbb{R} , its limit superior and limit inferior are

$$\limsup_{k \rightarrow \infty} a_k \equiv \begin{cases} \lim_{k \rightarrow \infty} \bar{a}_k & \text{if } \sup\{a_k\} < \infty, \\ \infty & \text{otherwise;} \end{cases}$$

$$\liminf_{k \rightarrow \infty} a_k \equiv \begin{cases} \lim_{k \rightarrow \infty} \underline{a}_k & \text{if } \inf\{a_k\} > -\infty, \\ -\infty & \text{otherwise.} \end{cases}$$

These are called simply the “lim sup” and “lim inf” for short.

Remark. By the Monotonic Sequence Theorem (Proposition 2.9) we have that

$$\limsup_{k \rightarrow \infty} a_k \equiv \begin{cases} \inf\{\bar{a}_k\} & \text{if } \sup\{a_k\} < \infty, \\ \infty & \text{otherwise;} \end{cases}$$

$$\liminf_{k \rightarrow \infty} a_k \equiv \begin{cases} \sup\{\underline{a}_k\} & \text{if } \inf\{a_k\} > -\infty, \\ -\infty & \text{otherwise.} \end{cases}$$

Example. Consider the sequence $\{a_k\}$ given by

$$a_k = (-1)^k \frac{k+1}{k} \quad \text{for } k \in \mathbb{Z}_+ .$$

The first eight terms of the sequences $\{\bar{a}_k\}$, $\{a_k\}$, and $\{\underline{a}_k\}$ are

| | | | | | | | | | |
|-------------------|--|---------------|----------------|----------------|----------------|----------------|----------------|----------------|-----------------|
| \bar{a}_k | | $\frac{3}{2}$ | $\frac{3}{2}$ | $\frac{5}{4}$ | $\frac{5}{4}$ | $\frac{7}{6}$ | $\frac{7}{6}$ | $\frac{9}{8}$ | $\frac{9}{8}$ |
| a_k | | -2 | $\frac{3}{2}$ | $-\frac{4}{3}$ | $\frac{5}{4}$ | $-\frac{6}{5}$ | $\frac{7}{6}$ | $-\frac{8}{7}$ | $\frac{9}{8}$ |
| \underline{a}_k | | -2 | $-\frac{4}{3}$ | $-\frac{4}{3}$ | $-\frac{6}{5}$ | $-\frac{6}{5}$ | $-\frac{8}{7}$ | $-\frac{8}{7}$ | $-\frac{10}{9}$ |

Notice that $\{a_k\}$ diverges while $\{\bar{a}_k\}$ and $\{\underline{a}_k\}$ are both monotonic and converge to 1 and -1 respectively. Therefore

$$\limsup_{k \rightarrow \infty} a_k = 1, \quad \liminf_{k \rightarrow \infty} a_k = -1 .$$

Remark. Notice that, unlike the limit, the lim sup and lim inf are defined for every real sequence, taking values in \mathbb{R}_{ex} , and that in general

$$-\infty \leq \liminf_{k \rightarrow \infty} a_k \leq \limsup_{k \rightarrow \infty} a_k \leq \infty .$$

Example. The sequence $\{(-1)^k k\}$ is neither bounded above nor bounded below. Therefore

$$\limsup_{k \rightarrow \infty} (-1)^k k = \infty, \quad \liminf_{k \rightarrow \infty} (-1)^k k = -\infty .$$

The key to learning how to use \limsup and \liminf is an understanding of the following characterizations. These should be compared with the characterization of the limit of a convergent sequence given by Proposition 2.6

Proposition 2.16. *Let $\{b_k\}$ be a sequence in \mathbb{R} . Let $b \in \mathbb{R}$. Then*

$$(2.13) \quad \limsup_{k \rightarrow \infty} b_k = b \quad \left(\liminf_{k \rightarrow \infty} b_k = b \right),$$

if and only if

$$(2.14) \quad b < c \quad \implies \quad b_k < c \quad \text{eventually (frequently)},$$

and

$$(2.15) \quad a < b \quad \implies \quad a < b_k \quad \text{frequently (eventually)}.$$

Proof. We give the proof of the \limsup assertion. The \liminf assertion is proved similarly.

(\implies) Suppose that (2.13) holds. Let $a < b$ and $c > b$. Then because

$$b = \limsup_{k \rightarrow \infty} b_k = \lim_{k \rightarrow \infty} \bar{b}_k,$$

where $\bar{b}_k = \sup\{b_l : l \geq k\}$, it follows that

$$a < \bar{b}_k < c \quad \text{eventually}.$$

Because $b_k \leq \bar{b}_k$ for every k , we see directly that $b_k < c$ eventually, whereby (2.14) holds. Moreover, $a < \bar{b}_k$ implies that for some $l \geq k$ one has $a < b_l$. (Otherwise a would be an upper bound for the set $\{b_l : l \geq k\}$, which contradicts the fact b is the least upper bound of this set.) Hence, $a < b_k$ frequently, whereby (2.15) holds.

(\impliedby) Suppose that (2.14) and (2.15) hold. Let $a < b$ and $c > b$ be arbitrary. Then (2.14) implies that

$$\bar{b}_k = \sup\{b_l : l \geq k\} \leq c \quad \text{eventually},$$

while (2.15) implies that

$$a < \bar{b}_k = \sup\{b_l : l \geq k\} \quad \text{eventually}.$$

(If for some k one had $\bar{b}_k \leq a$ then $b_l \leq a$ for every $l \geq k$, which contradicts (2.15).) Thus, we see

$$a \leq \inf\{\bar{b}_k\} \leq c.$$

But $a < b$ and $c > b$ were arbitrary, so that

$$\limsup_{k \rightarrow \infty} b_k = \inf\{\bar{b}_k\} = b,$$

whereby (2.13) holds. □

Convergent sequences may be characterized in terms of their \limsup and \liminf as follows.

Proposition 2.17. *Let $\{a_k\}$ be a sequence in \mathbb{R} . Then $\{a_k\}$ converges if and only if*

$$-\infty < \liminf_{k \rightarrow \infty} a_k = \limsup_{k \rightarrow \infty} a_k < \infty,$$

in which case

$$\lim_{k \rightarrow \infty} a_k = \liminf_{k \rightarrow \infty} a_k = \limsup_{k \rightarrow \infty} a_k.$$

Proof. Exercise. Hint: Use Propositions 2.6 and 2.16.

When adding and comparing sequences, \limsup and \liminf generally behave as follows.

Proposition 2.18. *Let $\{a_k\}$ and $\{b_k\}$ be sequences in \mathbb{R} . Then*

$$\begin{aligned}\limsup_{k \rightarrow \infty} (a_k + b_k) &\leq \limsup_{k \rightarrow \infty} a_k + \limsup_{k \rightarrow \infty} b_k, \\ \liminf_{k \rightarrow \infty} (a_k + b_k) &\geq \liminf_{k \rightarrow \infty} a_k + \liminf_{k \rightarrow \infty} b_k,\end{aligned}$$

whenever the sum on the right-hand side is defined. Moreover, if $a_k \leq b_k$ eventually then

$$\limsup_{k \rightarrow \infty} a_k \leq \limsup_{k \rightarrow \infty} b_k, \quad \liminf_{k \rightarrow \infty} a_k \leq \liminf_{k \rightarrow \infty} b_k.$$

Proof. Exercise.

When multiplying sequences, \limsup and \liminf generally behave as follows.

Proposition 2.19. *Let $\{a_k\}$ and $\{b_k\}$ be sequences in \mathbb{R} . If $\{a_k\}$ is convergent with*

$$\lim_{k \rightarrow \infty} a_k = a > 0,$$

then

$$\limsup_{k \rightarrow \infty} a_k b_k = a \limsup_{k \rightarrow \infty} b_k, \quad \liminf_{k \rightarrow \infty} a_k b_k = a \liminf_{k \rightarrow \infty} b_k.$$

Proof. Exercise.

When they converge, the \limsup and \liminf of a sequence are actually limits of some of its subsequences.

Proposition 2.20. *Let $\{a_k\}$ be a sequence in \mathbb{R} . If $\{a_{n_k}\}$ is any subsequence of $\{a_k\}$ then*

$$\liminf_{k \rightarrow \infty} a_k \leq \liminf_{k \rightarrow \infty} a_{n_k} \leq \limsup_{k \rightarrow \infty} a_{n_k} \leq \limsup_{k \rightarrow \infty} a_k.$$

Moreover, there exist subsequences $\{a_{n_k}\}$ and $\{a_{m_k}\}$ such that

$$\lim_{k \rightarrow \infty} a_{n_k} = \limsup_{k \rightarrow \infty} a_k, \quad \lim_{k \rightarrow \infty} a_{m_k} = \liminf_{k \rightarrow \infty} a_k.$$

Proof. Exercise. Hint: Use Propositions 2.6 and 2.16.

The following theorem associated with Bolzano and Weierstrass is an important consequence.

Proposition 2.21. (Bolzano-Weierstrass Theorem) *Every bounded sequence in \mathbb{R} has a convergent subsequence.*

Proof. Let $\{b_k\}$ be a bounded sequence in \mathbb{R} . This implies that there exists $[a, c] \subset \mathbb{R}$ such that $\{b_k\} \subset [a, c]$. Then

$$-\infty < a \leq \liminf_{k \rightarrow \infty} b_k \leq \limsup_{k \rightarrow \infty} b_k \leq c < \infty.$$

The result then follows by Proposition 2.20. □

2.8. Cauchy Criterion. When a sequence is monotonic, just knowing that it is bounded tells you that it is convergent. When a sequence is not monotonic, determining whether it is convergent or divergent is generally much harder. For example, to establish convergence directly from Definition 2.5 you must first know the limit of the sequence. Augustin Cauchy introduced a criterion for convergence that does not require knowledge of the limit.

Definition 2.9. A sequence $\{a_k\}_{k \in \mathbb{N}} \subset \mathbb{R}$ is said to be Cauchy whenever for every $\epsilon > 0$ there exists $N_\epsilon \in \mathbb{N}$ such that

$$(2.16) \quad k, l \geq N_\epsilon \implies |a_k - a_l| < \epsilon.$$

In other words, a sequence is Cauchy if for every $\epsilon > 0$ one can find a tail of the sequence such that any two terms in the tail are within ϵ of each other. Roughly speaking, a Cauchy sequence is one whose terms generally cluster together.

Remark. In order to show that a sequence is Cauchy, it is not enough to show merely that successive terms get closer together. Consider the sequence $\{a_k\}$ given by $a_k = \sqrt{k}$ for every $k \in \mathbb{N}$. Its successive terms get closer together because

$$\lim_{k \rightarrow \infty} (a_{k+1} - a_k) = \lim_{k \rightarrow \infty} (\sqrt{k+1} - \sqrt{k}) = \lim_{k \rightarrow \infty} \frac{1}{\sqrt{k+1} + \sqrt{k}} = 0.$$

However, $a_k \rightarrow \infty$ as $k \rightarrow \infty$, which implies $\{a_k\}$ is not Cauchy by Proposition 2.23 below.

The main result of this section is the so-called Cauchy criterion for convergence — namely, that a sequence in \mathbb{R} is convergent if and only if it is Cauchy. The easier half of this criterion is established by the following.

Proposition 2.22. A convergent sequence in \mathbb{R} is Cauchy.

Proof. Let $\{a_k\}$ be a convergent sequence in \mathbb{R} with limit a . Let $\epsilon > 0$. Then by the definition of convergence there exists $N_\epsilon \in \mathbb{N}$ such that

$$k \geq N_\epsilon \implies |a_k - a| < \frac{\epsilon}{2}.$$

It follows from the triangle inequality that if $k, l \geq N_\epsilon$ then

$$|a_k - a_l| = |(a_k - a) + (a - a_l)| \leq |a_k - a| + |a_l - a| < \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon.$$

Hence, the sequence $\{a_k\}$ is Cauchy. □

We now take the first step toward establishing the harder half of the Cauchy criterion.

Proposition 2.23. A Cauchy sequence in \mathbb{R} is bounded.

Proof. Let $\{a_k\}_{k \in \mathbb{N}}$ be a Cauchy sequence in \mathbb{R} . By Definition 2.9 there exists $N_1 \in \mathbb{N}$ such that

$$k, l \geq N_1 \implies |a_k - a_l| < 1.$$

This implies that for every $k \geq N_1$

$$|a_k| \leq |a_{N_1}| + |a_k - a_{N_1}| < |a_{N_1}| + 1.$$

Hence, for every $k \in \mathbb{N}$ we have

$$|a_k| < \max\{|a_l| : l = 0, \dots, N_1\} + 1.$$

The sequence $\{a_k\}_{k \in \mathbb{N}}$ is therefore bounded. □

We are now ready to establish the Cauchy Criterion.

Proposition 2.24. (Cauchy Criterion) *A sequence in \mathbb{R} is convergent if and only if it is Cauchy.*

Proof. Proposition 2.22 established that convergent sequences are Cauchy, so we only need to establish the other direction.

Let $\{a_k\}$ be a Cauchy sequence in \mathbb{R} . By Proposition 2.23 the sequence $\{a_k\}$ is bounded. By the Bolzano-Weierstrass Theorem (Proposition 2.21) it has a convergent subsequence $\{a_{n_k}\}$. Let a be the limit of this convergent subsequence. We will use the fact $\{a_k\}$ is a Cauchy sequence to show that it converges to a .

Let $\epsilon > 0$. Because the subsequence $\{a_{n_k}\}$ converges to a while the sequence $\{a_k\}$ is Cauchy, there exists an $N_\epsilon \in \mathbb{N}$ such that

$$k \geq N_\epsilon \implies |a_{n_k} - a| < \frac{\epsilon}{2},$$

and

$$k, l \geq N_\epsilon \implies |a_k - a_l| < \frac{\epsilon}{2}.$$

Because $k \geq N_\epsilon$ implies that $n_k \geq N_\epsilon$, the line above implies that

$$k \geq N_\epsilon \implies |a_k - a_{n_k}| < \frac{\epsilon}{2}.$$

It follows from the triangle inequality that if $k \geq N_\epsilon$ then

$$|a_k - a| \leq |a_k - a_{n_k}| + |a_{n_k} - a| < \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon.$$

The sequence $\{a_k\}$ therefore converges to a . □

2.9. Contracting Sequences* We now use the Cauchy Criterion to establish the convergence for a class of sequences that often arise in applications — namely, contracting sequences.

Definition 2.10. *A sequence $\{a_k\}_{k \in \mathbb{N}} \subset \mathbb{R}$ is said to be contracting if there exists $r \in (0, 1)$ such that*

$$(2.17) \quad |a_{k+1} - a_k| \leq r|a_k - a_{k-1}| \quad \text{eventually.}$$

In other words, a sequence is contracting if eventually the distance between successive points in it is reduced by at least a ratio $r < 1$.

Proposition 2.25. (Contracting Sequence) *Every contracting sequence in \mathbb{R} is convergent.*

Remark. Because Proposition 2.24 showed that Cauchy sequences are convergent, it suffices to show that contracting sequences are Cauchy.

Proof. Let $\{a_k\}_{k \in \mathbb{N}} \subset \mathbb{R}$ be contracting. Then there exists $r \in (0, 1)$ and $K \in \mathbb{N}$ such that (2.17) holds for every $k > K$. For every $k > K$ we can show by induction that

$$|a_k - a_{k+1}| < r|a_{k-1} - a_{k-2}| < r^2|a_{k-2} - a_{k-3}| < \cdots < r^{k-K}|a_K - a_{K+1}|.$$

The triangle inequality and the difference of powers formula then imply that for $l > k > K$ we have the bound

$$\begin{aligned} |a_k - a_l| &\leq |a_k - a_{k+1}| + |a_{k+1} - a_{k+2}| + \cdots + |a_{l-2} - a_{l-1}| + |a_{l-1} - a_l| \\ &< r^{k-K} (1 + r + \cdots + r^{l-k-2} + r^{l-k-1}) |a_K - a_{K+1}| \\ &= r^{k-K} \frac{1 - r^{l-k}}{1 - r} |a_k - a_{k+1}| < \frac{r^{k-K}}{1 - r} |a_K - a_{K+1}|. \end{aligned}$$

Now let $\epsilon > 0$. Because $r < 1$ we can pick $N_\epsilon > K$ such that $r^{N_\epsilon - K} |a_K - a_{K+1}| < (1 - r)\epsilon$. Let $k, l > N_\epsilon$ with $k \neq l$. Without loss of generality we may assume that $k < l$. By the above bound we have

$$|a_k - a_l| < \frac{r^{k-K}}{1-r} |a_K - a_{K+1}| < \frac{r^{N_\epsilon - K}}{1-r} |a_K - a_{K+1}| < \epsilon.$$

Therefore $\{a_k\}_{k \in \mathbb{N}}$ is Cauchy, whereby it is convergent by Proposition 2.24. \square

Remark. The next exercise shows that not every convergent sequence is contracting.

Exercise. Show that $\{1/k\}_{k \in \mathbb{Z}_+}$ is not contracting.

Remark. The next example illustrates one way to show a sequence is contracting. We will see other ways later in the course.

Example. Let $a_0 > 0$ and $a_{k+1} = 1/(2 + a_k^2)$ for every $k \in \mathbb{N}$. Show $\{a_k\}_{k \in \mathbb{N}}$ is contracting.

We first express $a_{k+1} - a_k$ in terms of a_k and a_{k-1} by using the defining relation twice as

$$\begin{aligned} a_{k+1} - a_k &= \frac{1}{2 + a_k^2} - \frac{1}{2 + a_{k-1}^2} = \frac{a_{k-1}^2 - a_k^2}{(2 + a_k^2)(2 + a_{k-1}^2)} = \frac{a_{k-1} + a_k}{(2 + a_k^2)(2 + a_{k-1}^2)} (a_{k-1} - a_k) \\ &= \left(\frac{a_{k-1}}{2 + a_{k-1}^2} \frac{1}{2 + a_k^2} + \frac{1}{2 + a_{k-1}^2} \frac{a_k}{2 + a_k^2} \right) (a_{k-1} - a_k). \end{aligned}$$

Next, use the fact that $a_k \in (0, \frac{1}{2})$ for every $k \geq 1$ and the elementary inequalities

$$\frac{x}{2 + x^2} \leq \frac{2}{9}, \quad \frac{1}{2 + x^2} < \frac{1}{2}, \quad \text{for every } x \in (0, \frac{1}{2}),$$

to derive the bound

$$|a_{k+1} - a_k| < \left(\frac{2}{9} \frac{1}{2} + \frac{1}{2} \frac{2}{9} \right) |a_k - a_{k-1}| = \frac{2}{9} |a_k - a_{k-1}| \quad \text{for every } k \geq 2.$$

This shows that $\{a_k\}_{k \in \mathbb{N}}$ is contracting. It is also convergent by Proposition 2.25. \square

Exercise. Let $a_0 > 0$ and $a_{k+1} = 1/(2 + a_k)$ for every $k \in \mathbb{N}$. Show $\{a_k\}_{k \in \mathbb{N}}$ is contracting.

Exercise. Let $a_0 > 1$ and $a_{k+1} = 1/(1 + a_k)$ for every $k \in \mathbb{N}$. Show $\{a_k\}_{k \in \mathbb{N}}$ is contracting.

Exercise. Let $a_0 > 1$ and $a_{k+1} = 1/(1 + a_k^2)$ for every $k \in \mathbb{N}$. Show $\{a_k\}_{k \in \mathbb{N}}$ is contracting.

3. SERIES OF REAL NUMBERS

3.1. Infinite Series. Any finite set of real numbers can be summed. Here we study one way to make sense of the sum of an infinite sequence of real numbers.

Definition 3.1. Given any real sequence $\{a_k\}_{k=0}^{\infty}$, for every $m, n \in \mathbb{N}$ with $m \leq n$ define the sigma notation:

$$\sum_{k=m}^n a_k \equiv a_m + a_{m+1} + \cdots + a_{n-1} + a_n.$$

Associate with the sequence of terms $\{a_k\}$ the so-called sequence of partial sums $\{s_n\}$ defined by

$$s_n \equiv \sum_{k=0}^n a_k.$$

It is convenient to encode $\{s_n\}$ with the formal infinite series

$$\sum_{k=0}^{\infty} a_k.$$

If the sequence $\{s_n\}$ converges to a limit s then we say that the series converges, and that s is the sum of the series. In that case we write

$$\sum_{k=0}^{\infty} a_k = s.$$

If the sequence $\{s_n\}$ diverges then we say that the series diverges.

Remark. It is clear that changing, adding, or removing a finite number of terms in a series does not affect whether the series converges or diverges, but if it converges, the sum would almost always be affected. For example,

$$\sum_{k=0}^{\infty} a_k \text{ converges} \iff \sum_{k=5}^{\infty} a_k \text{ converges},$$

but when they do converge the sums will generally differ — namely,

$$\text{in general } \sum_{k=0}^{\infty} a_k \neq \sum_{k=5}^{\infty} a_k.$$

More specifically, these sums will be equal if and only if $a_0 + a_1 + a_2 + a_3 + a_4 = 0$.

Example. Consider the infinite series

$$\sum_{k=1}^{\infty} \frac{1}{k(k+1)}.$$

The n^{th} partial sum is given by

$$\begin{aligned} s_n &= \sum_{k=1}^n \frac{1}{k(k+1)} = \sum_{k=1}^n \left(\frac{1}{k} - \frac{1}{k+1} \right) \\ &= \left(1 - \frac{1}{2} \right) + \left(\frac{1}{2} - \frac{1}{3} \right) + \cdots + \left(\frac{1}{n} - \frac{1}{n+1} \right) \\ &= 1 + \left(-\frac{1}{2} + \frac{1}{2} \right) + \cdots + \left(-\frac{1}{n} + \frac{1}{n} \right) - \frac{1}{n+1} = 1 - \frac{1}{n+1}. \end{aligned}$$

One then sees that the series is convergent with

$$\sum_{k=1}^{\infty} \frac{1}{k(k+1)} = \lim_{n \rightarrow \infty} s_n = 1.$$

The previous example shows a series that can be put into so-called *telescoping form*.

Definition 3.2. A formal infinite series with terms $\{a_k\}_{k=m}^{\infty}$ is said to be in telescoping form if $a_k = c_{k-1} - c_k$ for some sequence $\{c_k\}_{k=m-1}^{\infty}$, so that the series is expressed as

$$(3.1) \quad \sum_{k=m}^{\infty} (c_{k-1} - c_k).$$

If a series can be put into telescoping form with a sequence $\{c_k\}$ that is known explicitly then the convergence or divergence of the series can be easily determined. Moreover, if it converges then its sum can be easily determined. This is because the sequence $\{c_k\}$ is simply related to the sequence of partial sums $\{s_n\}$. Indeed, for every $n \geq m$ one sees that

$$\begin{aligned} s_n &= \sum_{k=m}^n a_k = \sum_{k=m}^n (c_{k-1} - c_k) \\ &= (c_{m-1} - c_m) + (c_m - c_{m+1}) + \cdots + (c_{n-2} - c_{n-1}) + (c_{n-1} - c_n) \\ &= c_{m-1} + (-c_m + c_m) + \cdots + (-c_{n-1} + c_{n-1}) - c_n \\ &= c_{m-1} - c_n. \end{aligned}$$

It follows immediately that the sequence $\{s_n\}$ converges if and only if the sequence $\{c_n\}$ converges, and that when these sequences converge one has that

$$\lim_{n \rightarrow \infty} s_n = c_{m-1} - \lim_{n \rightarrow \infty} c_n.$$

Hence, the following proposition holds.

Proposition 3.1. Let $\{c_k\}_{k=m-1}^{\infty}$ be a real sequence. Then

$$\sum_{k=m}^{\infty} (c_{k-1} - c_k) \text{ converges} \quad \iff \quad \{c_k\}_{k=m-1}^{\infty} \text{ converges}.$$

Moreover, when these are convergent one has

$$\sum_{k=m}^{\infty} (c_{k-1} - c_k) = c_{m-1} - \lim_{k \rightarrow \infty} c_k.$$

Remark. The above considerations show that if a series is in the telescoping form (3.1) then there is a $c \in \mathbb{R}$ such that $c_k = c - s_k$ for every $k \geq m$, where $\{s_k\}_{k=m}^{\infty}$ is the sequence of partial sums. This means that finding an explicit telescoping form for a series is equivalent to finding an explicit expression for its partial sums. It should be clear that one can do this in only the rarest of cases.

Given a general infinite series, it is usually impossible to evaluate the limit of the sequence of partial sums. However, one can commonly determine whether a series is convergent or divergent without explicitly evaluating this limit. The following proposition gives the simplest test for divergence.

Proposition 3.2. (Divergence Test) *Let $\{a_k\}$ be a real sequence.*

If the series $\sum_{k=0}^{\infty} a_k$ converges then $\lim_{k \rightarrow \infty} a_k = 0$.

Equivalently, if $\lim_{k \rightarrow \infty} a_k \neq 0$ then the series $\sum_{k=0}^{\infty} a_k$ diverges.

Proof. The proof is based on the fact that the k^{th} term in a formal infinite series can be expressed as $a_k = s_k - s_{k-1}$, where $s_{-1} = 0$ and $\{s_k\}_{k \in \mathbb{N}}$ is the sequence of partial sums. If the series converges then one knows that

$$\lim_{k \rightarrow \infty} s_k = s, \quad \lim_{k \rightarrow \infty} s_{k-1} = s,$$

where s is the sum of the series. It thereby follows that

$$\lim_{k \rightarrow \infty} a_k = \lim_{k \rightarrow \infty} s_k - \lim_{k \rightarrow \infty} s_{k-1} = s - s = 0. \quad \square$$

Remark. You can easily find examples of a series whose terms converge to zero, yet the series is divergent. One such example is the *harmonic series*:

$$\sum_{k=1}^{\infty} \frac{1}{k}.$$

Clearly $1/k \rightarrow 0$ as $k \rightarrow \infty$. However, we will soon show that this series diverges.

3.2. Geometric Series. An important example is that of geometric series.

Definition 3.3. *A formal infinite series of the form*

$$\sum_{k=0}^{\infty} ar^k$$

for some nonzero a and some $r \in \mathbb{R}$ is called a geometric series.

The convergence or divergence of a geometric series is easy to determine because it is one of those rare series where one can find an explicit expression for its partial sums. For every $n \in \mathbb{N}$ let s_n denote the partial sum given by

$$s_n = \sum_{k=0}^n ar^k.$$

It is clear that if $r = 1$ then $s_n = (n + 1)a$ and the series will diverge. So suppose that $r \neq 1$. One checks that

$$s_n - rs_n = \sum_{k=0}^n ar^k - \sum_{k=0}^n ar^{k+1} = \sum_{k=0}^n ar^k - \sum_{k=1}^{n+1} ar^k = a - ar^{n+1},$$

whereby the partial sum s_n is found to be

$$s_n = \frac{a - ar^{n+1}}{1 - r}.$$

By letting n tend to ∞ in this expression we find that

$$\sum_{k=0}^{\infty} ar^k = \begin{cases} \frac{a}{1 - r} & \text{if } |r| < 1, \\ \text{diverges} & \text{otherwise.} \end{cases}$$

Remark. The fact the geometric series diverges when $|r| \geq 1$ can also be seen easily from the Divergence Test. Indeed, in that case you see that

$$\lim_{k \rightarrow \infty} ar^k = \begin{cases} a & \text{(and hence is nonzero) if } r = 1, \\ \text{diverges} & \text{(and hence is nonzero) if } |r| \geq 1 \text{ and } r \neq 1, \end{cases}$$

whereby the Divergence Test (Proposition 3.2) shows that the geometric series diverges. Of course, the Divergence Test does not show the geometric series converges when $|r| < 1$.

Exercise. Consider a formal infinite series of the form

$$\sum_{k=1}^{\infty} kr^k$$

for some $r \in \mathbb{R}$. Find all the values of r for which this series converges and evaluate the sum. (Hint: Find an explicit expression for the partial sums and evaluate the limit. The explicit expression may be derived from the analogous expression for a geometric series.)

3.3. Series with Nonnegative Terms. If the terms of an infinite series are nonnegative then the associated sequence of partial sums will be nondecreasing. Hence, the least upper bound property can be employed in the guise of the Monotonic Sequence Theorem (Proposition 2.9) to show the convergence or divergence of the series. Specifically, one has the following proposition, which lies at the heart of most proofs about the convergence or divergence of series with nonnegative terms.

Proposition 3.3. (Series with Nonnegative Terms Theorem) *Let $\{a_k\}_{k=m}^{\infty}$ be a nonnegative sequence. Then*

$$\sum_{k=m}^{\infty} a_k \text{ converges} \iff \{s_k\}_{k=m}^{\infty} \text{ is bounded above,}$$

where $\{s_k\}_{k=m}^{\infty}$ is the sequence of partial sums associated with the formal infinite series.

Proof. One first shows that the sequence $\{s_k\}_{k=m}^{\infty}$ is nondecreasing. One then applies the Monotonic Sequence Theorem. The details are left as an exercise. \square

One way to establish whether or not a sequence of partial sums is bounded above is to compare it with a sequence of partial sums for which the answer is known. This is often done with one of the following comparison tests.

Proposition 3.4. (Comparison Tests for Series with Nonnegative Terms) *Let $\{a_k\}$ and $\{b_k\}$ be nonnegative sequences that satisfy one of the following comparison conditions:*

(i) *the direct comparison*

$$\exists M \in \mathbb{R}_+ \quad \text{such that} \quad a_k \leq M b_k \quad \text{eventually};$$

(ii) *the limit comparison (if each b_k is positive)*

$$\limsup_{k \rightarrow \infty} \frac{a_k}{b_k} < \infty;$$

(iii) *the ratio comparison (if each a_k and b_k is positive)*

$$\frac{a_{k+1}}{a_k} \leq \frac{b_{k+1}}{b_k} < \infty \quad \text{eventually.}$$

Then

$$(3.2) \quad \sum_{k=0}^{\infty} b_k \quad \text{converges} \quad \Longrightarrow \quad \sum_{k=0}^{\infty} a_k \quad \text{converges,}$$

$$\left(\sum_{k=0}^{\infty} a_k \quad \text{diverges} \quad \Longrightarrow \quad \sum_{k=0}^{\infty} b_k \quad \text{diverges.} \right)$$

Proof. First, condition (i) implies (3.2) because if $a_k \leq M b_k$ for every $k \geq m$ then the fact that $\sum b_k$ converges yields the upper bound

$$\sum_{k=m}^n a_k \leq M \sum_{k=m}^n b_k \leq M \sum_{k=m}^{\infty} b_k < \infty.$$

Proposition 3.3 therefore implies that $\sum a_k$ converges. Next, condition (ii) implies condition (i) (and hence (3.2)) upon observing that

$$\limsup_{k \rightarrow \infty} \frac{a_k}{b_k} < M < \infty \quad \Longrightarrow \quad a_k \leq M b_k \quad \text{eventually.}$$

Finally, condition (iii) implies condition (ii) (and hence (3.2)) upon observing that

$$\frac{a_{k+1}}{a_k} \leq \frac{b_{k+1}}{b_k} < \infty \quad \text{eventually} \quad \Longrightarrow \quad \frac{a_{k+1}}{b_{k+1}} \leq \frac{a_k}{b_k} \quad \text{eventually}$$

$$\Longrightarrow \quad \left\{ \frac{a_k}{b_k} \right\} \quad \text{is nonincreasing eventually}$$

$$\Longrightarrow \quad \lim_{k \rightarrow \infty} \frac{a_k}{b_k} < \infty. \quad \square$$

Exercise. The proof argues that the direct comparison test applies whenever the limit comparison test applies, and that the limit comparison test applies whenever the ratio comparison test applies. Can you find an examples where (a) the direct comparison test applies but the limit comparison test fails, or (b) the limit comparison test applies but the ratio comparison test fails?

Example. We can apply the direct comparison test to show the harmonic series diverges. Consider the comparison

$$1 \leq 1, \quad \frac{1}{2} \leq \frac{1}{2}, \quad \frac{1}{4} \leq \frac{1}{3}, \quad \frac{1}{4} \leq \frac{1}{4}, \quad \frac{1}{8} \leq \frac{1}{5}, \quad \frac{1}{8} \leq \frac{1}{6}, \quad \frac{1}{8} \leq \frac{1}{7}, \quad \frac{1}{8} \leq \frac{1}{8}, \quad \dots$$

Summing both sides, one sees that

$$1 + \frac{n}{2} \leq \sum_{k=1}^{2^n} \frac{1}{k}.$$

The partial sums clearly diverge.

3.4. A Series for e . Recall that we defined e to be the real number such that

$$(3.3) \quad \lim_{n \rightarrow \infty} \left(1 + \frac{1}{n}\right)^n = \lim_{n \rightarrow \infty} \left(1 - \frac{1}{n}\right)^{-n} = e.$$

Here we will show that e is also the sum of an infinite series. What makes this example different from the previous ones is that here we will evaluate the sum without passing to the limit in an explicit expression for the partial sums. Rather, we will compare the partial sums with first sequence in (3.3).

Proposition 3.5. *The number e satisfies*

$$(3.4) \quad \sum_{k=0}^{\infty} \frac{1}{k!} = e.$$

Remark. This result is due to Euler. Newton had showed that the solution $y = \exp(x)$ of the initial-value problem $y' = y$, $y(0) = 1$ behaved like an exponential function and was given by

$$\exp(x) = \sum_{k=0}^{\infty} \frac{x^k}{k!}.$$

Its base, $\exp(1)$ is given by the series in (3.4). Euler showed this base is e as defined by (3.3).

Proof. Define the sequences $\{a_n\}_{n=1}^{\infty}$ and $\{s_n\}_{n=0}^{\infty}$ by

$$a_n = \left(1 + \frac{1}{n}\right)^n, \quad s_n = \sum_{k=0}^n \frac{1}{k!}.$$

We want to use the fact that $a_n \rightarrow e$ to show that $s_n \rightarrow e$. The result will then follow because $\{s_n\}_{n=0}^{\infty}$ is the sequence of partial sums associated with the infinite series in (3.4).

For $n \geq 2$ the binomial formula yields

$$a_n = \left(1 + \frac{1}{n}\right)^n = \sum_{k=0}^n \frac{n!}{k!(n-k)!} \frac{1}{n^k} = 1 + 1 + \sum_{k=2}^n \frac{1}{k!} \prod_{l=1}^{k-1} \left(1 - \frac{l}{n}\right).$$

Because the product in each term of the last sum is bounded above by 1, we see that $a_n \leq s_n$. Hence, the sequence of partial sums satisfies

$$(3.5) \quad e = \lim_{n \rightarrow \infty} a_n \leq \liminf_{n \rightarrow \infty} s_n.$$

Let $m \in \mathbb{Z}_+$. For every $n > m$ we have

$$a_n \geq \sum_{k=0}^m \frac{n!}{k!(n-k)!} \frac{1}{n^k} = 1 + 1 + \sum_{k=2}^m \frac{1}{k!} \prod_{l=1}^{k-1} \left(1 - \frac{l}{n}\right).$$

Letting $n \rightarrow \infty$ in this inequality while keeping m fixed shows that s_m is bounded above as

$$e = \lim_{n \rightarrow \infty} a_n \geq s_m.$$

Because the sequence $\{s_n\}$ is increasing and bounded above by e , by the Monotone Sequence Theorem (Proposition 2.9) it converges with

$$(3.6) \quad \lim_{m \rightarrow \infty} s_m \leq e.$$

This upper bound together with the lower bound (3.5) shows that $s_n \rightarrow e$ as $n \rightarrow \infty$, and thereby establishes (3.4). \square

We can use this result to prove that e is not rational.

Proposition 3.6. *The number e is irrational.*

Proof. Suppose e is rational. Then $e = p/q$ for some $p, q \in \mathbb{Z}_+$. By Proposition 3.5

$$e = \sum_{k=0}^{\infty} \frac{1}{k!}$$

Let

$$s_q = \sum_{k=0}^q \frac{1}{k!}.$$

Then by subtracting s_q from e and comparing the resulting infinite series with a geometric series we see that

$$\begin{aligned} 0 < e - s_q &= \sum_{k=q+1}^{\infty} \frac{1}{k!} = \frac{1}{(q+1)!} + \frac{1}{(q+2)!} + \frac{1}{(q+3)!} + \cdots \\ &< \frac{1}{(q+1)!} \left(1 + \frac{1}{q+1} + \frac{1}{(q+1)^2} + \cdots \right) = \frac{1}{q!q}. \end{aligned}$$

Upon multiplying these inequalities by $q!$ we arrive at

$$0 < q!(e - s_q) < \frac{1}{q}.$$

But $q!e = (q-1)!p$ is an integer while $q!s_q$ is also an integer because

$$q!s_q = \sum_{k=0}^q \frac{q!}{k!}.$$

Therefore $q!(e - s_q)$ is an integer that lies in $(0, \frac{1}{q})$, which is a contradiction because $q \geq 1$. \square

Remark. In fact, e is transcendental. A real number is said to be *algebraic* if it is a zero of some polynomial with integer coefficients. Real numbers that are not algebraic are said to be *transcendental*.

3.5. Series with Nonincreasing Positive Terms. The harmonic series is a special case of the so-called p -series, which is formally given by

$$(3.7) \quad \sum_{k=1}^{\infty} \frac{1}{k^p}.$$

Because the terms of this series are nonincreasing and positive, the following two convergence tests can be applied. Proposition 3.3 plays a central role in their proofs.

Proposition 3.7. (Cauchy 2^k Test) *Let $\{a_k\}$ be a nonincreasing, positive sequence. Then*

$$\sum_{k=1}^{\infty} a_k \text{ converges} \iff \sum_{k=0}^{\infty} 2^k a_{2^k} \text{ converges}.$$

Proof. The result is a consequence of the direct comparisons

$$a_{2^{j+1}} \leq a_k \leq a_{2^j} \quad \text{for } 2^j \leq k < 2^{j+1},$$

which yield the bounds

$$\sum_{j=0}^{n-1} 2^{j-1} a_{2^{j+1}} \leq \sum_{k=1}^{2^n-1} a_k \leq \sum_{j=0}^{n-1} 2^j a_{2^j} \quad \text{for every } n \in \mathbb{Z}_+.$$

The details are left as an exercise. □

Example. Because $\{1/k^p\}$ is a nonincreasing, positive sequence, Proposition 3.7 implies that the p -series (3.7) converges or diverges as the series

$$\sum_{k=0}^{\infty} 2^k \frac{1}{2^{kp}} = \sum_{k=0}^{\infty} (2^{1-p})^k.$$

But this is a geometric series that clearly converges for $p > 1$ and diverges for $p \leq 1$.

Remark. The proof of the Cauchy 2^k test outlined above extends the argument by which we showed the harmonic series diverges. Indeed, the harmonic series is just the p -series for $p = 1$.

Remark. The Cauchy 2^k test can be generalized to subsequences of $\{a_k\}$ of the form $\{a_{n_k}\}$ where there exist constants \underline{m} and \overline{m} such that

$$0 < \underline{m} \leq \frac{n_{k+1} - n_k}{n_k - n_{k-1}} \leq \overline{m} < \infty.$$

For example, one can choose $n_k = m^k$ for some $m \in \mathbb{N}$ with $m > 1$. This satisfies

$$\frac{n_{k+1} - n_k}{n_k - n_{k-1}} = \frac{m - 1}{1 - m^{-1}} = m,$$

whereby $\underline{m} = \overline{m} = m$. This leads to a “ m^k test” Its statement and proof is left to you.

The second convergence test of this section requires the use of integrals — in fact, the use of *improper* integrals. These will be developed rigorously later in the course. However, here we will assume you have some familiarity with them from your elementary calculus courses.

Proposition 3.8. (Integral Test) *Let f be a nonincreasing, positive, locally integrable (continuous, for example) function over $[0, \infty)$. Then*

$$\sum_{k=0}^{\infty} f(k) \text{ converges} \iff \int_0^{\infty} f(x) dx \text{ converges}.$$

Remark. The integral above is understood in the sense of an improper integral.

Proof. The key fact we need from integration theory is that the improper integral

$$\int_0^{\infty} f(x) \, dx \quad \text{converges}$$

whenever the sequence $\{S_n\}$ converges, where each S_n is defined by

$$S_n = \int_0^n f(x) \, dx.$$

Because $\{S_n\}$ is an increasing sequence, showing convergence reduces to showing it is bounded above.

The result will then be a consequence of the fact that

$$S_n = \sum_{k=1}^n \int_{k-1}^k f(x) \, dx \quad \text{for every } n \in \mathbb{Z}_+,$$

and the direct comparisons

$$f(k) \leq \int_{k-1}^k f(x) \, dx \leq f(k-1), \quad \text{for every } k \in \mathbb{Z}_+.$$

These facts should be clear to you based on your knowledge of definite integrals from elementary calculus. If not, a picture should help clarify things. We will establish them rigorously later. Here we will assume they are true and complete the proof.

By summing the above direct comparisons, one obtains

$$\sum_{k=1}^n f(k) \leq S_n \leq \sum_{k=1}^n f(k-1) = \sum_{k=0}^{n-1} f(k).$$

The remaining details are left as an exercise. □

Example. Because $\{1/k^p\}$ is a nonincreasing, positive sequence, Proposition 3.8 implies that the p -series (3.7) converges or diverges as the improper integral

$$\int_1^{\infty} \frac{1}{x^p} \, dx.$$

But for $p \neq 1$ one can easily check that

$$S_k = \int_1^k \frac{1}{x^p} \, dx = \frac{1}{p-1} \left(1 - \frac{1}{k^{p-1}} \right),$$

while for $p = 1$ one has

$$S_k = \int_1^k \frac{1}{x} \, dx = \log(k).$$

(Here $\log(\cdot)$ denotes the natural logarithm.) One then sees that the sequence $\{S_k\}$ converges for $p > 1$ and diverges for $p \leq 1$. The same is therefore true for the p -series.

3.6. Alternating Series. Until now we have only studied convergence tests for nonnegative series. The underlying tool has been the Monotonic Sequence Theorem (Proposition 2.9), which was used to prove Proposition 3.3. Here we use the Monotonic Sequence Theorem to obtain the following characterization of convergence for a special class of series with alternating sign.

Proposition 3.9. (Alternating Series Test) *Let $\{a_k\}$ be a positive, nonincreasing sequence in \mathbb{R} . Then*

$$\sum_{k=0}^{\infty} (-1)^k a_k \text{ converges} \iff \lim_{k \rightarrow \infty} a_k = 0.$$

Proof. The direction \implies is just Proposition 3.2 (Divergence Test). To prove the other direction, let

$$s_n = \sum_{k=0}^n (-1)^k a_k.$$

First, the picture is that

$$\{s_{2k}\}_{k \in \mathbb{N}} \text{ is nonincreasing, while } \{s_{2k+1}\}_{k \in \mathbb{N}} \text{ is nondecreasing,}$$

and that

$$s_{2k} > s_{2j+1} \text{ for every } j, k \in \mathbb{N}.$$

Indeed, the first two assertions follow because

$$\begin{aligned} s_{2k+2} - s_{2k} &= a_{2k+2} - a_{2k+1} \leq 0, \\ s_{2k+3} - s_{2k+1} &= -a_{2k+3} + a_{2k+2} \geq 0. \end{aligned}$$

Next, because $s_{2k} > s_{2k+1}$ for every $k \in \mathbb{N}$, for any $j \leq k$ one has

$$s_{2k} > s_{2k+1} \geq s_{2j+1}, \quad s_{2j} \geq s_{2k} > s_{2k+1}.$$

The result follows by exchanging j and k in the last inequality. The monotonic subsequences $\{s_{2k}\}$ and $\{s_{2k+1}\}$ are thereby bounded below and above respectively. By the Monotonic Sequence Theorem they therefore converge. Let

$$\bar{s} = \lim_{k \rightarrow \infty} s_{2k}, \quad \underline{s} = \lim_{k \rightarrow \infty} s_{2k+1}.$$

Then

$$\bar{s} - \underline{s} = \lim_{k \rightarrow \infty} (s_{2k} - s_{2k+1}) = \lim_{k \rightarrow \infty} a_{2k+1} = 0,$$

whereby $\bar{s} = \underline{s}$. The last step is to show that this fact implies that $\{s_k\}$ converges. This is left as an exercise. \square

Examples.

$$\sum_{k=1}^{\infty} \frac{(-1)^k}{k^p} \text{ converges for } p > 0, \quad \sum_{k=2}^{\infty} \frac{(-1)^k}{\log(k)} \text{ converges.}$$

3.7. Absolute Convergence. The Monotonic Sequence Theorem has been the tool underlying all the convergence tests we have studied so far. We now use the Cauchy criterion to establish a test that does not require the series to be nonnegative.

Proposition 3.10. (Absolute Convergence Test) *Let $\{a_k\}$ be a real sequence. Then*

$$\sum_{k=0}^{\infty} |a_k| \text{ converges} \implies \sum_{k=0}^{\infty} a_k \text{ converges.}$$

Proof. Let $\{p_n\}$ and $\{q_n\}$ be the sequences of partial sums given by

$$p_n = \sum_{k=0}^n |a_k|, \quad q_n = \sum_{k=0}^n a_k.$$

By hypotheses $\{p_n\}$ is convergent, and thereby Cauchy. The idea of the proof is to show that $\{q_n\}$ is Cauchy, and thereby convergent.

The key to doing so is the fact that for every $m, n \in \mathbb{N}$ one has the inequality

$$|q_n - q_m| \leq |p_n - p_m|.$$

This is trivially true when $m = n$. When $n > m$ the triangle inequality yields

$$|q_n - q_m| = \left| \sum_{k=m+1}^n a_k \right| \leq \sum_{k=m+1}^n |a_k| = |p_n - p_m|.$$

The case $n < m$ goes similarly.

Let $\epsilon > 0$. Because $\{p_n\}$ is Cauchy there exists an $N_\epsilon \in \mathbb{N}$ such that

$$m, n \geq N_\epsilon \implies |p_n - p_m| < \epsilon.$$

Because $|q_n - q_m| \leq |p_n - p_m|$, one immediately sees that

$$m, n \geq N_\epsilon \implies |q_n - q_m| < \epsilon.$$

Hence, $\{q_n\}$ is Cauchy, and thereby convergent. □

Proposition 3.10 motivates the following definition.

Definition 3.4. *If $\{a_k\}$ is a real sequence such that*

$$\sum_{k=0}^{\infty} |a_k| \text{ converges,}$$

then one says

$$\sum_{k=0}^{\infty} a_k \text{ converges absolutely,}$$

or

$$\text{is absolutely convergent.}$$

Example. Consider the alternating p -series

$$\sum_{n=1}^{\infty} \frac{(-1)^n}{n^p}.$$

This converges for $p > 0$ by the alternating series test, but it converges absolutely only for $p > 1$. This example shows that not every convergent series is absolutely convergent. In other words, absolute convergence is a stronger property than convergence.

When the definition of absolute convergence is combined with the Comparison Tests for series with nonnegative terms (Proposition 3.4), we get an array of new comparison tests for absolute convergence that can be applied to general series.

Proposition 3.11. (Absolute Comparison Tests) *Let $\{a_k\}$ and $\{b_k\}$ be real sequences that satisfy one of the following comparison conditions:*

(i) *the direct comparison*

$$\exists M \in \mathbb{R}_+ \quad \text{such that} \quad |a_k| \leq M b_k \quad \text{eventually};$$

(ii) *the limit comparison (if each b_k is positive)*

$$\limsup_{k \rightarrow \infty} \frac{|a_k|}{b_k} < \infty;$$

(iii) *the ratio comparison (if each $|a_k|$ and b_k is positive)*

$$\frac{|a_{k+1}|}{|a_k|} \leq \frac{b_{k+1}}{b_k} < \infty \quad \text{eventually}.$$

Then

$$(3.8) \quad \sum_{k=0}^{\infty} b_k \quad \text{converges} \quad \implies \quad \sum_{k=0}^{\infty} a_k \quad \text{converges absolutely}.$$

Proof. Exercise.

Example. Because $|\cos(kx)| \leq 1$ for every $x \in \mathbb{R}$ and $k \in \mathbb{Z}_+$, direct comparison with the p -series shows that the series

$$\sum_{k=1}^{\infty} \frac{\cos(kx)}{k^p} \quad \text{converges absolutely for } p > 1.$$

3.8. Root and Ratio Tests. The root and ratio tests both draw their conclusions about the convergence of a series based on absolute comparisons with a geometric series.

Proposition 3.12. (Root Test) *Let $\{a_k\}$ be a real sequence. Let*

$$\rho = \limsup_{k \rightarrow \infty} \sqrt[k]{|a_k|}.$$

Then

$$\begin{aligned} \rho < 1 &\implies \sum_{k=0}^{\infty} a_k \quad \text{converges absolutely,} \\ \rho > 1 &\implies \sum_{k=0}^{\infty} a_k \quad \text{diverges.} \end{aligned}$$

If $\rho = 1$ the series may either converge or diverge.

Proof. The convergence conclusion when $\rho < 1$ follows by a direct comparison of the series with a convergent geometric series. Specifically, by Proposition 2.16 one has that

$$\rho < r < 1 \implies \limsup_{k \rightarrow \infty} \sqrt[k]{|a_k|} < r \implies |a_k| < r^k \quad \text{eventually.}$$

The absolute convergence follows from the direct comparison test of Proposition 3.11.

The divergence conclusion when $\rho > 1$ follows by showing that $\limsup |a_k| > 0$. Specifically,

$$1 < r < \rho \implies \limsup_{k \rightarrow \infty} \sqrt[k]{|a_k|} > r \implies |a_k| > r^k \text{ frequently.}$$

This implies there exists a subsequence $\{a_{n_k}\}_k$ of $\{a_k\}_k$ such that

$$|a_{n_k}| > r^{n_k} \text{ eventually.}$$

Then $\limsup |a_k| \geq \limsup |a_{n_k}| \geq \lim r^{n_k} = \infty$. But $\limsup |a_k| > 0$ implies the sequence $\{a_k\}$ does not converge to zero, which by the Divergence Test (Proposition 3.2) implies the associated series diverges.

We leave as an exercise the task of finding examples of both a convergent and a divergent series with $\rho = 1$. \square

Proposition 3.13. (Ratio Test) *Let $\{a_k\}$ be a nonzero real sequence. Then*

$$\begin{aligned} \limsup_{k \rightarrow \infty} \frac{|a_{k+1}|}{|a_k|} < 1 &\implies \sum_{k=0}^{\infty} a_k \text{ converges absolutely,} \\ \frac{|a_{k+1}|}{|a_k|} \geq 1 \text{ eventually} &\implies \sum_{k=0}^{\infty} a_k \text{ diverges.} \end{aligned}$$

Proof. As with the proof of the root test, the convergence conclusion follows by a direct comparison of the series with a convergent geometric series, while the divergence conclusion follows by showing that $\limsup |a_k| > 0$. Specifically, by Proposition 2.16 one has that

$$\rho < r < 1 \implies \limsup_{k \rightarrow \infty} \frac{|a_{k+1}|}{|a_k|} < r \implies \frac{|a_{k+1}|}{|a_k|} < r \text{ eventually.}$$

An induction argument can then be used to show that for some $m \in \mathbb{N}$ one has

$$|a_k| \leq |a_m| r^{k-m} \text{ for every } k \geq m.$$

Because the geometric series

$$\sum_{k=m}^{\infty} \frac{|a_m|}{r^m} r^k \text{ converges,}$$

the comparison theorem implies

$$\sum_{k=0}^{\infty} a_k \text{ converges.}$$

The proof of the divergence assertion is left as an exercise. \square

Remark. Some books give the divergence criterion of the ratio test as

$$\liminf_{k \rightarrow \infty} \frac{|a_{k+1}|}{|a_k|} > 1 \implies \sum_{k=0}^{\infty} a_k \text{ diverges.}$$

This is clearly weaker than the one we give.

Remark. The version of the ratio test most commonly found in elementary calculus texts makes a much stronger hypothesis than the above version. It is the following.

Proposition 3.14. (Elementary Ratio Test) Let $\{a_k\}$ be a nonzero real sequence such that

$$(3.9) \quad \rho = \lim_{k \rightarrow \infty} \frac{|a_{k+1}|}{|a_k|} \quad \text{exists.}$$

Then

$$\begin{aligned} \rho < 1 &\implies \sum_{k=0}^{\infty} a_k \quad \text{converges absolutely,} \\ \rho > 1 &\implies \sum_{k=0}^{\infty} a_k \quad \text{diverges.} \end{aligned}$$

If $\rho = 1$ the series may either converge or diverge.

Hypothesis (3.9) above requires the existence of a limit, whereas Proposition 3.13 has no such requirement. (Recall that the lim sup exists in \mathbb{R}_{ex} for every sequence, even ones that diverge.)

Example. Find the least upper bound and greatest lower bound of the set

$$S = \left\{ x \in \mathbb{R} : \sum_{n=0}^{\infty} \frac{(3n)!}{n!} \frac{(2n)!}{(4n)!} x^n \quad \text{converges} \right\}.$$

This can be easily done employing the ratio test. Indeed, because

$$\frac{|a_{n+1}|}{|a_n|} = \frac{\frac{(3n+3)!}{(n+1)!} \frac{(2n+2)!}{(4n+4)!} |x|^{n+1}}{\frac{(3n)!}{n!} \frac{(2n)!}{(4n)!} |x|^n} = \frac{(3n+3)(3n+2)(3n+1)(2n+2)(2n+1)}{(n+1)(4n+4)(4n+3)(4n+2)(4n+1)} |x|.$$

one finds that

$$\lim_{n \rightarrow \infty} \frac{|a_{n+1}|}{|a_n|} = \frac{3^3 \times 2^2}{4^4} |x| = \frac{3^3}{4^3} |x|.$$

The elementary ratio test, Proposition 3.14, implies that the series converges when $|x| < (4/3)^3$ and diverges when $|x| > (4/3)^3$. Therefore $\sup\{S\} = (4/3)^3$ while $\inf\{S\} = -(4/3)^3$. The root test can be applied with the aid of the factorial-root limit (2.12) to obtain the same result.

Remark. The root test is sometimes harder to apply, but as the following indicates, its convergence assertion can be sharper.

Proposition 3.15. Let $\{a_k\}$ be a positive sequence. Then

$$(3.10) \quad \liminf_{k \rightarrow \infty} \frac{a_{k+1}}{a_k} \leq \liminf_{k \rightarrow \infty} \sqrt[k]{a_k} \leq \limsup_{k \rightarrow \infty} \sqrt[k]{a_k} \leq \limsup_{k \rightarrow \infty} \frac{a_{k+1}}{a_k}.$$

Proof. Exercise. (The middle inequality is obvious, so just prove the other two.)

Exercise. Find a series for which all the inequalities in (3.10) are strict.

Remark. Because both the root and ratio tests draw their convergence conclusions based on comparison with a geometric series, they should only be used when such a comparison makes sense. For example, these tests can be used to assert the absolute convergence of series like

$$\sum_{k=1}^{\infty} k^4 2^{-k}, \quad \sum_{l=0}^{\infty} e^{-l^2} 4^l, \quad \sum_{m=0}^{\infty} \frac{(m!)^2}{(2m)!} (-3)^m,$$

but will yield no information about the convergence of series like

$$\sum_{k=2}^{\infty} \frac{\log(k)}{k^2}, \quad \sum_{l=0}^{\infty} \left(\frac{3l+2}{l^4+2} \right)^{\frac{1}{2}}, \quad \sum_{m=2}^{\infty} \frac{(-1)^m}{m(\log(m))^2}.$$

3.9. Dirichlet Test*. We now apply the Cauchy criterion to establish a test that, like the Alternating Series Test, can be applied to series that do not converge absolutely.

Proposition 3.16. (Dirichlet Test) *Let $\{a_k\}_{k \in \mathbb{N}}$ be a positive, nonincreasing sequence in \mathbb{R} such that*

$$\lim_{k \rightarrow \infty} a_k = 0.$$

Let $\{b_k\}_{k \in \mathbb{N}}$ be a sequence in \mathbb{R} for which there exists M such that

$$(3.11) \quad \left| \sum_{k=0}^n b_k \right| \leq M \quad \text{for every } n \in \mathbb{N}.$$

Then

$$\sum_{k=0}^{\infty} a_k b_k \quad \text{converges.}$$

Remark. The Dirichlet Test implies the convergence conclusion of the Alternating Series Test. Indeed, if we set $b_k = (-1)^k$ then

$$\sum_{k=0}^n b_k = \sum_{k=0}^n (-1)^k = \begin{cases} 1 & \text{for } n \text{ even,} \\ 0 & \text{for } n \text{ odd.} \end{cases}$$

Hence, bound (3.11) holds with $M = 1$. The Dirichlet Test then tells us that for every positive, nonincreasing real sequence $\{a_k\}_{k \in \mathbb{N}}$ such that $a_k \rightarrow 0$ as $k \rightarrow \infty$ the series

$$\sum_{k=0}^{\infty} (-1)^k a_k \quad \text{converges.}$$

As the next example illustrates, the Dirichlet Test is far more powerful than the Alternating Series Test.

Example. Consider the problem of determining all $x, p \in \mathbb{R}$ for which the Fourier p -series

$$\sum_{k=1}^{\infty} \frac{\cos(kx)}{k^p} \quad \text{converges.}$$

Because $|\cos(kx)| \leq 1$ for every $k \in \mathbb{Z}_+$, direct comparison with the regular p -series shows that this series converges absolutely for $p > 1$. However this argument says nothing about what happens when $p \leq 1$.

First observe that when $x \in \{2m\pi : m \in \mathbb{Z}\}$ one has $\cos(kx) = 1$ for every $k \in \mathbb{Z}_+$. In this case the Fourier p -series reduces to a regular p -series, which diverges for every $p \leq 1$.

Next, observe that when $x \in \{(2m+1)\pi : m \in \mathbb{Z}\}$ one has $\cos(kx) = (-1)^k$ for every $k \in \mathbb{Z}_+$. In this case the Fourier p -series reduces to an alternating p -series, which (by the Alternating Series Test) converges for every $p > 0$.

We now use the Dirichlet Test to analyze the more general case when $x \notin \{2m\pi : m \in \mathbb{Z}\}$. Let $a_k = 1/k^p$ and $b_k = \cos(kx)$. Clearly the sequence $\{a_k\}$ is positive, decreasing, and vanishes

as $k \rightarrow \infty$. The hard step is to show that the partial sums associated with the sequence $\{b_k\}$ satisfy (3.11). To do this we use the trigonometric identity

$$2 \sin\left(\frac{1}{2}x\right) \cos(kx) = \sin\left(\left(k + \frac{1}{2}\right)x\right) - \sin\left(\left(k - \frac{1}{2}\right)x\right),$$

and the fact $\sin\left(\frac{1}{2}x\right) \neq 0$ when $x \notin \{2m\pi : m \in \mathbb{Z}\}$ to obtain (by a telescoping sum) the formula

$$\sum_{k=1}^n b_k = \sum_{k=1}^n \cos(kx) = \sum_{k=1}^n \frac{\sin\left(\left(k + \frac{1}{2}\right)x\right) - \sin\left(\left(k - \frac{1}{2}\right)x\right)}{2 \sin\left(\frac{1}{2}x\right)} = \frac{\sin\left(\left(n + \frac{1}{2}\right)x\right) - \sin\left(\frac{1}{2}x\right)}{2 \sin\left(\frac{1}{2}x\right)}.$$

It is clear from this formula that

$$\left| \sum_{k=1}^n b_k \right| = \left| \frac{\sin\left(\left(n + \frac{1}{2}\right)x\right) - \sin\left(\frac{1}{2}x\right)}{2 \sin\left(\frac{1}{2}x\right)} \right| \leq \frac{|\sin\left(\left(n + \frac{1}{2}\right)x\right)| + |\sin\left(\frac{1}{2}x\right)|}{2 |\sin\left(\frac{1}{2}x\right)|} \leq \frac{1}{|\sin\left(\frac{1}{2}x\right)|}.$$

Hence, bound (3.11) holds with $M = 1/|\sin\left(\frac{1}{2}x\right)|$. The Dirichlet Test then implies that when $x \notin \{2m\pi : m \in \mathbb{Z}\}$ the Fourier p -series converges for every $p > 0$.

Finally, you can use the Divergence Test to show that the Fourier p -series diverges for every $p \leq 0$. This follows easily once you know that

$$\limsup_{k \rightarrow \infty} \cos(kx) > 0 \quad \text{for every } x \in \mathbb{R}.$$

We leave the details as an exercise. □

Remark. When applying the Dirichlet test to a given series, one must identify the sequences $\{a_k\}$ and $\{b_k\}$, and check that all the hypotheses on them are satisfied. The hypotheses on $\{a_k\}$ are easy to check, so do that first: the sequence $\{a_k\}$ must be positive, nonincreasing, and vanish as $k \rightarrow \infty$. The hypothesis on $\{b_k\}$ is typically much harder to check: the associated partial sums must satisfy (3.11). The key to checking this in the above example was to write $b_k = c_{k+1} - c_k$ (by using a trigonometric identity) for some bounded sequence $\{c_k\}$, whereby the partial sums telescoped as

$$\sum_{k=0}^n b_k = \sum_{k=0}^n (c_{k+1} - c_k) = c_{n+1} - c_0.$$

This telescoping approach can be taken for a variety of other $\{b_k\}$ too.

Our proof of the Dirichlet Test uses an identity that is a discrete analog of the integration-by-parts formula from calculus. Because it has many other applications, this identity gets its own proposition.

Proposition 3.17. (Summation-by-Parts Identity) *Let $\{a_k\}_{k \in \mathbb{N}}$ and $\{b_k\}_{k \in \mathbb{N}}$ be sequences in \mathbb{R} . Let $B_{-1} = 0$ and*

$$B_n = \sum_{k=0}^n b_k \quad \text{for every } n \in \mathbb{N}.$$

Then for every $m, n \in \mathbb{N}$ with $m \leq n$ one has identity

$$\sum_{k=m}^n a_k b_k = a_n B_n - a_m B_{m-1} + \sum_{k=m}^{n-1} (a_k - a_{k+1}) B_k,$$

with the understanding that the last sum is zero when $m = n$.

Remark. This is called the summation-by-parts identity because it is a discrete analog of the integration-by-parts formula

$$\int_m^n a(x)b(x) \, dx = a(x)B(x) \Big|_m^n - \int_m^n a'(x)B(x) \, dx,$$

where $B'(x) = b(x)$.

Proof. Because $b_k = B_k - B_{k-1}$ we have

$$\begin{aligned} \sum_{k=m}^n a_k b_k &= \sum_{k=m}^n a_k (B_k - B_{k-1}) \\ &= \sum_{k=m}^n a_k B_k - \sum_{k=m}^n a_k B_{k-1} \\ &= \sum_{k=m}^n a_k B_k - \sum_{k=m-1}^{n-1} a_{k+1} B_k \\ &= a_n B_n - a_m B_{m-1} + \sum_{k=m}^{n-1} (a_k - a_{k+1}) B_k. \end{aligned}$$

To get from the second to the third line above we re-indexed the last sum. All the other steps are straightforward algebra. \square

We now turn to the proof of the Dirichlet Test.

Proof of Dirichlet Test. Let

$$s_n = \sum_{k=0}^n a_k b_k.$$

We will show the sequence $\{s_k\}_{k \in \mathbb{N}}$ is Cauchy, and therefore convergent.

Let $\epsilon > 0$. We seek $N_\epsilon \in \mathbb{N}$ such that

$$m, n \geq N_\epsilon \implies |s_n - s_m| < \epsilon.$$

For $m = n$ this is always true. Suppose $m < n$. (For the case $n < m$ simply reverse the roles of m and n .) Then

$$\begin{aligned} |s_n - s_m| &= \left| \sum_{k=m+1}^n a_k b_k \right| = \left| a_n B_n - a_{m+1} B_m + \sum_{k=m+1}^{n-1} (a_k - a_{k+1}) B_k \right| \\ &\leq a_n |B_n| + a_{m+1} |B_m| + \sum_{k=m+1}^{n-1} (a_k - a_{k+1}) |B_k| \\ &\leq a_n M + a_{m+1} M + \sum_{k=m+1}^{n-1} (a_k - a_{k+1}) M = 2a_{m+1} M. \end{aligned}$$

Here we have used the summation-by-parts identity in the second step, the triangle inequality and the fact that $\{a_k\}_{k \in \mathbb{N}}$ is positive and nonincreasing in the third step, the bound $|B_k| \leq M$ in the fourth step, and evaluated the telescoping sum in the last step. Because $a_k \rightarrow 0$ as $k \rightarrow \infty$, we can choose N_ϵ so that $m \geq N_\epsilon$ implies $2a_{m+1} M < \epsilon$. Hence, for every $n > m \geq N_\epsilon$ the above inequalities imply $|s_n - s_m| < \epsilon$. \square

4. SETS OF REAL NUMBERS

4.1. Closure, Closed, and Dense. The notions of *closure*, *closed*, and *dense* pertain to sets as they relate to the limit process. The closure of a set is all points that are the limit of some convergent sequence that lies within the set. If the closure of a set is the set itself then the set is said to be closed. Simply put, limits do not get out of closed sets. If the closure of a set is everything then the set is said to be dense. Simply put, limits can go anywhere from dense sets. Here we make these notions precise for subsets of \mathbb{R} .

4.1.1. *Closure.* We begin with the definition.

Definition 4.1. Given any $A \subset \mathbb{R}$ its closure is given by

$$A^c = \left\{ a \in \mathbb{R} : a \text{ is the limit of a sequence in } A \right\}.$$

It is clear that $A \subset A^c$ for every $A \subset \mathbb{R}$. Indeed, every $a \in A$ is the limit of the constant sequence $\{a_k\}$ with $a_k = a$ for every $k \in \mathbb{N}$. As we will now see, sometimes $A^c = A$, but in general A^c will be larger than A .

Examples. It is easy to show that $\emptyset^c = \emptyset$ and $\mathbb{R}^c = \mathbb{R}$.

Examples. If $a < b$ then the closures of the intervals (a, b) , $(a, b]$, $[a, b)$, $[a, b]$, (a, ∞) , $[a, \infty)$, $(-\infty, b)$, and $(-\infty, b]$ are given by

$$\begin{aligned} (a, b)^c &= (a, b]^c = [a, b)^c = [a, b]^c = [a, b], \\ (a, \infty)^c &= [a, \infty)^c = [a, \infty), \quad (-\infty, b)^c = (-\infty, b]^c = (-\infty, b]. \end{aligned}$$

You should be able to prove these facts.

We have the following propositions.

Proposition 4.1. If $A \subset \mathbb{R}$ is nonempty and bounded above (below) then $\sup\{A\} \in A^c$ ($\inf\{A\} \in A^c$).

Proof. You have to show that if $\sup\{A\} \notin A$ then there exists a sequence $\{a_k\} \subset A$ such that $a_k \rightarrow \sup\{A\}$ as $k \rightarrow \infty$. The details are left as an exercise. \square

Proposition 4.2. For every $A, B \subset \mathbb{R}$ one has that

- (i) $A \subset B \implies A^c \subset B^c$,
- (ii) $(A \cup B)^c = A^c \cap B^c$,
- (iii) $(A \cap B)^c \supset A^c \cup B^c$.

Proof. Exercise.

An important fact is that the closure of \mathbb{Q} is \mathbb{R} . In other words, every real number is the limit of a sequence of rational numbers.

Proposition 4.3. $\mathbb{Q}^c = \mathbb{R}$.

Proof. Let $a \in \mathbb{R}$. Consider the sequence of intervals $\{I_k\}_{k \in \mathbb{N}}$ where each I_k is given by

$$I_k = \left(a - \frac{1}{2^k}, a + \frac{1}{2^k} \right).$$

For each $k \in \mathbb{N}$ the third assertion of Proposition 1.15 implies there exists a $a_k \in I_k \cap \mathbb{Q}$. The step of showing that $a_k \rightarrow a$ as $k \rightarrow \infty$ is left as an exercise. It then follows that $a \in \mathbb{Q}^c$, whereby the assertion follows. \square

4.1.2. *Closed.* We are ready for the next definition.

Definition 4.2. A subset A of \mathbb{R} is said to be closed when $A = A^c$.

Examples. The empty set \emptyset is closed.

Examples. If $a < b$ then intervals of the form $[a, a]$, $[a, b]$, $[a, \infty)$, $(-\infty, b]$, and $\mathbb{R} = (-\infty, \infty)$ are closed, while intervals of the form (a, b) , $(a, b]$, $[a, b)$, (a, ∞) , and $(-\infty, b)$ are not.

Remark. Earlier we had used the word “closed” to describe those intervals in \mathbb{R} that contain their endpoints. The foregoing examples show that our new usage of “closed” coincides with that old usage. The new usage can also apply to sets that are not intervals, so is more general.

Our terminology seems to demand that closures should be closed. This is indeed the case.

Proposition 4.4. Let $A \subset \mathbb{R}$. Then A^c is closed (i.e. $(A^c)^c = A^c$).

Proof. Let $a \in (A^c)^c$. We must show that $a \in A^c$. Because $a \in (A^c)^c$ there exists a sequence $\{b_i\}_{i \in \mathbb{N}}$ in A^c such that $b_i \rightarrow a$ as $i \rightarrow \infty$. If $b_i = a$ for some $i \in \mathbb{N}$ then $a = b_i \in A^c$.

On the other hand, if $b_i \neq a$ for every $i \in \mathbb{N}$ then because $b_i \in A^c$ for each $i \in \mathbb{N}$ there exists a sequence $\{b_{(i,j)}\}_{j \in \mathbb{N}}$ in A such that $b_{(i,j)} \rightarrow b_i$ as $j \rightarrow \infty$. The picture is

$$\begin{array}{ccccccc}
 b_{(0,0)}, & b_{(0,1)}, & b_{(0,2)}, & \cdots & b_{(0,j)}, & \cdots & \rightarrow & b_0 \\
 b_{(1,0)}, & b_{(1,1)}, & b_{(1,2)}, & \cdots & b_{(1,j)}, & \cdots & \rightarrow & b_1 \\
 b_{(2,0)}, & b_{(2,1)}, & b_{(2,2)}, & \cdots & b_{(2,j)}, & \cdots & \rightarrow & b_2 \\
 \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
 b_{(i,0)}, & b_{(i,1)}, & b_{(i,2)}, & \cdots & b_{(i,j)}, & \cdots & \rightarrow & b_i \\
 \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
 & & & & & & & \downarrow \\
 & & & & & & & a,
 \end{array}$$

Because for each $i \in \mathbb{N}$ we have $b_{(i,j)} \rightarrow b_i$ as $j \rightarrow \infty$ and $|b_i - a| > 0$, there exists a $j_i \in \mathbb{N}$ such that

$$|b_{(i,j_i)} - b_i| < |b_i - a|.$$

Set $a_i = b_{(i,j_i)}$ for each $i \in \mathbb{N}$. It is clear that the sequence $\{a_i\}_{i \in \mathbb{N}}$ lies within A . The step of showing that $a_i \rightarrow a$ as $i \rightarrow \infty$ is left as an exercise. It then follows that $a \in A^c$, whereby the assertion follows. \square

Proposition 4.5. Let $A \subset \mathbb{R}$. Then A^c is the smallest closed set that contains A .

Proof. The previous proposition shows that A^c is closed. Earlier we showed that $A \subset A^c$. Now let B be any closed set that contains A . We see from (i) of Proposition 4.2 that $A \subset B$ implies $A^c \subset B^c$. Because B is closed we know that $B^c = B$. It follows that $A^c \subset B^c = B$. Therefore A^c is the smallest closed set that contains A . \square

The property of being closed is preserved by certain set operations.

Proposition 4.6. If A and B are closed subsets of \mathbb{R} then $A \cap B$ and $A \cup B$ are closed. If $\{A_k\}_{k \in \mathbb{N}}$ is a sequence of closed subsets of \mathbb{R} then

$$\bigcap_{k \in \mathbb{N}} A_k \quad \text{is closed.}$$

Proof. Exercise.

Remark. By repeated application of the first assertion above, we see that the union and intersection of any finite collection of closed sets is again closed. The second assertion states that the intersection of any countable collection of closed sets is again closed. The analogous statement for unions is generally false. Indeed, consider the countable collection of closed intervals $\{I_k\}_{k \in \mathbb{N}}$ where each I_k is given by

$$I_k = \left[-1 + \frac{1}{2^k}, 1 - \frac{1}{2^k} \right].$$

You can easily show that

$$\bigcup_{k \in \mathbb{N}} I_k = (-1, 1),$$

which is not closed.

4.1.3. *Dense.* Finally, we have the concept of a set being dense in a larger one.

Definition 4.3. Let $A \subset B \subset \mathbb{R}$. Then A is said to be dense in B if $B \subset A^c$.

Examples. Proposition 4.3 states the \mathbb{Q} is dense in \mathbb{R} . In a similar manner one can show that $(a, b) \cap \mathbb{Q}$ is dense in $[a, b]$, that $(a, \infty) \cap \mathbb{Q}$ is dense in $[a, \infty)$, and $(-\infty, b) \cap \mathbb{Q}$ is dense in $(-\infty, b]$.

Proposition 4.7. If $A \subset B \subset C \subset D \subset \mathbb{R}$ and A is dense in D then B is dense in C .

Proof. Exercise.

Proposition 4.8. Let $A \subset \mathbb{R}$. Then A is dense in \mathbb{R} if and only if for every interval (a, b) one has $A \cap (a, b) \neq \emptyset$.

Proof. Exercise.

4.2. **Completeness.** Completeness is a central notion regarding sets in analysis. As such, it arises in many settings. Here we introduce it in the setting of subsets of \mathbb{R} , where it is easily characterized. The basic notion of a set being complete is as follows.

Definition 4.4. A set $S \subset \mathbb{R}$ is said to be complete if every Cauchy sequence contained in S has a limit that is in S .

The Cauchy Criterion, Proposition 2.24, immediately implies that \mathbb{R} is complete. Moreover, it easily yields the following characterization of all complete subsets of \mathbb{R} .

Proposition 4.9. A subset of \mathbb{R} is complete if and only if it is closed.

Proof. Exercise.

Remark. In more general settings the notions of complete and closed do not coincide. For example, consider the set \mathbb{Q} equipped with the usual notion of distance. A sequence in \mathbb{Q} is said to be convergent in \mathbb{Q} if it is convergent as a sequence in \mathbb{R} and its limit is in \mathbb{Q} . A sequence in \mathbb{Q} is said to be Cauchy in \mathbb{Q} if it is Cauchy as a sequence in \mathbb{R} . Because there are sequences in \mathbb{Q} that are Cauchy in \mathbb{Q} but not convergent in \mathbb{Q} the set \mathbb{Q} is not complete. On the other hand, the set \mathbb{Q} is closed because it contains all possible limit points of sequences that are convergent in \mathbb{Q} .

4.3. Connectedness*. Connectedness is another central notion regarding sets in analysis. As such, it comes in many varieties. Fortunately, these varieties coincide in the setting of subsets of \mathbb{R} . The basic notion of a set being connected is as follows.

Definition 4.5. *A set $S \subset \mathbb{R}$ is said to be disconnected there exists nonempty $A, B \subset S$ such that*

$$(4.1) \quad A \cup B = S, \quad A^c \cap B = A \cap B^c = \emptyset.$$

Otherwise S is said to be connected.

If a set S is disconnected then the nonempty sets A and B that arise in Definition 4.5 have the property that any convergent sequence that lies within one them will have a limit that is not in the other. In other words, if $\{x_n\} \subset A$ is convergent and $x_n \rightarrow x$ then $x \notin B$.

Example. The set $(-\infty, 0) \cup (0, \infty)$ is disconnected because (4.1) is satisfied by the sets $A = (-\infty, 0)$ and $B = (0, \infty)$.

It is clear from Definition 4.5 that every disconnected set has at least two points in it. In particular, the empty set or any singleton set (a set containing only a single point) is connected. The following proposition shows that the connected subsets of \mathbb{R} are precisely the intervals. Our proof will use the Interval Characterization Theorem, Proposition 1.19.

Proposition 4.10. *A subset of \mathbb{R} is connected if and only if it is an interval.*

Remark. This shows that Definition 4.5 of a connected set leads to a result in accord with what one might naively expect connectedness to mean for subsets of \mathbb{R} .

Proof. (\implies) Let $S \subset \mathbb{R}$ be connected. We will show that S is then an interval by using the Interval Characterization Theorem, Proposition 1.19.

Let $x, y \in S$ such that $x < y$. We must show that $(x, y) \subset S$. Let $z \in (x, y)$. We must show that $z \in S$. Suppose not. Let $A = (-\infty, z] \cap S$ and $B = [z, \infty) \cap S$. Because $x \in A$ and $y \in B$, these sets are nonempty. It is easy to check that A and B satisfy (4.1), whereby S is disconnected. But this contradicts the fact S is connected. Hence, $z \in S$. Therefore $(x, y) \subset S$. Because this is true for every $x, y \in S$ such that $x < y$, the Interval Characterization Theorem, Proposition 1.19, implies that S is an interval.

(\impliedby) Let $S \subset \mathbb{R}$ be an interval. Suppose that S is disconnected. Then there exists nonempty sets $A, B \subset S$ that satisfy (4.1). Let $x \in A$ and $y \in B$. Because we can always relabel the sets A and B , we can assume without loss of generality that $x < y$. Because $x, y \in S$ while S is an interval, we know from the Interval Characterization Theorem, Proposition 1.19, that $[x, y] \subset S$. Because $A \cup B = S$, we have $(A \cap [x, y]) \cup (B \cap [x, y]) = [x, y]$.

Now consider the point $z = \sup\{A \cap [x, y]\}$. By Propositions 4.1 and 4.2 we have

$$z \in (A \cap [x, y])^c \subset A^c \cap [x, y].$$

Because $A^c \cap B = \emptyset$ and $y \in B$, it follows that $z \neq y$, which implies that $z < y$. Because $z = \sup\{A \cap [x, y]\} < y$ while $(A \cap [x, y]) \cup (B \cap [x, y]) = [x, y]$, we see that $(z, y) \subset B$. But then $z \in [z, y] \subset B^c$ by Proposition 4.2, so that $z \notin A$ because $A \cap B^c = \emptyset$. On the other hand, because $A^c \cap B = \emptyset$ while $(A \cap [x, y]) \cup (B \cap [x, y]) = [x, y]$, it follows that $z \in A \cap [x, y] \subset A$. This contradicts the conclusion of the sentence before it. Therefore S must be connected. \square

4.4. Sequential Compactness. Compactness is a central notion regarding sets in analysis that plays a crucial role in many existence proofs. As such, it also comes in many varieties. Fortunately, these varieties coincide in the setting of subsets of \mathbb{R} . We will take advantage of this coincidence by presenting only the concept of sequential compactness, for which we have all the tools at hand.

Definition 4.6. *A set $A \subset \mathbb{R}$ is said to be sequentially compact if every sequence in A has a subsequence that converges to a limit in A .*

Example. The interval $[0, \infty)$ is not sequentially compact because the increasing sequence $\{k\}_{k \in \mathbb{N}}$ diverges to ∞ , and therefore has no convergent subsequence.

Example. The interval $(0, 1)$ is not sequentially compact because the limit of the convergent sequence $\{2^{-k}\}_{k \in \mathbb{N}}$ is 0, which is not in $(0, 1)$.

4.4.1. Characterization. It is clear that every sequentially compact set must be closed, for otherwise there would be a convergent sequence within it whose limit lies outside it. Intuitively, a sequentially compact set must also be “small enough” that every sequence within it has a convergent subsequence. The following characterization of sequentially compact subsets of \mathbb{R} uses the Bolzano-Weierstrass Theorem to show that “small enough” is simply that the set is bounded.

Proposition 4.11. *A set $A \subset \mathbb{R}$ is sequentially compact if and only if A is closed and bounded.*

Proof. (\implies) Suppose that A is either not bounded or not closed.

If A is not bounded then for every $k \in \mathbb{N}$ there exists $a_k \in A$ such that $|a_k| > k$. But then the sequence $\{|a_k|\}$ diverges to ∞ , whereby every subsequence of $\{|a_k|\}$ diverges to ∞ . But then Propositions 2.7 and 2.8 imply that every subsequence of $\{a_k\}$ diverges. Hence, $\{a_k\}$ is a sequence in A that has no convergent subsequence. Therefore A is not sequentially compact.

If A is not closed then there exists a sequence $\{a_k\}$ in A and a point $a \notin A$ such that $a_k \rightarrow a$ as $k \rightarrow \infty$. By Proposition 2.7 every subsequence of $\{a_k\}$ also converges to the point a , which is not in A . Therefore A is not sequentially compact.

(\impliedby) Let $A \subset \mathbb{R}$ be closed and bounded. Let $\{a_k\}$ be an arbitrary sequence in A . Because A is bounded, the sequence $\{a_k\}$ is bounded. By the Bolzano-Weierstrass Theorem, $\{a_k\}$ has a converging subsequence $\{a_{n_k}\}$. Let a be the limit of this subsequence. Because A is closed and $\{a_{n_k}\}$ is in A , the limit a must also be in A . By the arbitrariness of $\{a_k\}$, we conclude that every sequence in A has a subsequence that converges to a limit in A . Therefore A is sequentially compact. \square

An immediate consequence of this characterization is the following.

Proposition 4.12. *Let $A \subset \mathbb{R}$ be sequentially compact and $B \subset \mathbb{R}$ be closed. Then $A \cap B$ is sequentially compact. In particular, every closed subset of A is sequentially compact.*

Proof. Exercise.

It also follows that the property of being sequentially compact is preserved by unions.

Proposition 4.13. *If A and B are sequentially compact subsets of \mathbb{R} then so is $A \cup B$.*

Proof. Exercise.

Remark. By repeated application of this proposition, we see that the union of any finite collection of sequentially compact sets is again sequentially compact.

4.4.2. *Open Interval Covering Property.* Sequentially compact subsets of \mathbb{R} have many covering properties, the simplest of which is the following.

Proposition 4.14. *Let $A \subset \mathbb{R}$ be sequentially compact. Let $\{(a_k, b_k)\}_{k \in \mathbb{N}}$ be a countable collection of open intervals that covers A — i.e. such that*

$$A \subset \bigcup_{k \in \mathbb{N}} (a_k, b_k).$$

Then there exist $n \in \mathbb{N}$ such that

$$A \subset \bigcup_{k=0}^n (a_k, b_k).$$

Remark. This property is often stated as: every countable covering of A by open intervals has a finite subcovering. Many other covering properties have a similar flavor — namely, that every covering of A by a certain class of sets has a smaller subcovering.

Proof. Suppose not. Then for every $n \in \mathbb{N}$ there exists an $x_n \in A$ such that

$$(4.2) \quad x_n \notin \bigcup_{k=0}^n (a_k, b_k).$$

Because $\{x_n\}_{n \in \mathbb{N}} \subset A$ while A is sequentially compact, there exists a subsequence $\{x_{n_k}\}_{k \in \mathbb{N}}$ that converges to a limit $x \in A$. Because $\{(a_k, b_k)\}_{k \in \mathbb{N}}$ covers A , there exists $m \in \mathbb{N}$ such that $x \in (a_m, b_m)$. Because $x_{n_k} \rightarrow x$ as $k \rightarrow \infty$, this implies that $x_{n_k} \in (a_m, b_m)$ eventually as $k \rightarrow \infty$. But this contradicts the fact seen from (4.2) that $x_{n_k} \notin (a_m, b_m)$ for every $n_k \geq m$. \square

There is a converse of the previous proposition that we state without proof.

Proposition 4.15. *Let $A \subset \mathbb{R}$ such that for every countable collection of open intervals $\{(a_k, b_k)\}_{k \in \mathbb{N}}$ that covers A there exist $n \in \mathbb{N}$ such that*

$$A \subset \bigcup_{k=0}^n (a_k, b_k).$$

Then A is sequentially compact.

Remark. By combining Propositions 4.14 and 4.15, we see that the covering property stated in the hypothesis of Proposition 4.15 characterizes sequential compactness for subsets of \mathbb{R} . This property is closely related to the properties of *countable compactness* and *compactness*, which are also covering properties that characterize sequential compactness for subsets of \mathbb{R} . In more general settings these notions of compactness can differ from each other.

4.4.3. *Cantor Intersection Theorem.* Sequential compactness lies behind the following useful generalization by Cantor of the Nested Interval Theorem.

Proposition 4.16. (Cantor Intersection Theorem) *Let $\{A_n\}_{n \in \mathbb{N}}$ be a nested countable collection of nonempty, closed, bounded subsets of \mathbb{R} — i.e. nonempty, closed, bounded sets such that*

$$A_0 \supset A_1 \supset \cdots \supset A_n \supset A_{n+1} \supset \cdots .$$

Then

$$\bigcap_{n \in \mathbb{N}} A_n \neq \emptyset .$$

Proof. Because each A_n is nonempty, for every $n \in \mathbb{N}$ there exists $x_n \in A_n$. Consider the sequence $\{x_n\}$. Because the collection $\{A_n\}$ is nested, for every $m, n \in \mathbb{N}$ such that $n \geq m$ one has $A_n \subset A_m$, whereby $x_n \in A_m$. The sequence $\{x_n\}$ is thereby in each A_m eventually.

Because A_0 is closed and bounded, it is sequentially compact by Proposition 4.11. Because A_0 is sequentially compact while $\{x_n\} \subset A_0$, there exists a subsequence $\{x_{n_k}\}_{k \in \mathbb{N}}$ that converges to a limit $x \in A_0$. Because the subsequence $\{x_{n_k}\}$ is in each A_m eventually while each A_m is closed, it follows that x is in each A_m . Hence, $x \in \bigcap_{m \in \mathbb{N}} A_m$, whereby the intersection is nonempty. \square

Remark. The hypothesis that the A_n are bounded is essential. For example, consider the nested collection of nonempty, closed sets $\{[n, \infty)\}_{n \in \mathbb{N}}$. Clearly,

$$\bigcap_{n \in \mathbb{N}} [n, \infty) = \emptyset.$$

The next proposition will explore this further.

Proposition 4.16 provides half of the following characterization of sequentially compactness.

Proposition 4.17. *Let $A \subset \mathbb{R}$ be closed. Then A is sequentially compact if and only if every countable collection of nested, nonempty, closed subsets of A has a nonempty intersection.*

Proof. (\implies) This follows from Propositions 4.11 and 4.16.

(\impliedby) Suppose A is not sequentially compact. Then there exists a sequence $\{x_n\}_{n \in \mathbb{N}} \subset A$ such that no subsequence of it converges to a point in A . We claim that no subsequence of $\{x_n\}$ converges. Indeed, because A is closed, any subsequence of $\{x_n\}$ that converges must converge to a point in A , which is a contradiction. For each $m \in \mathbb{N}$ define $B_m = \{x_n : n \geq m\}$. We leave as an exercise the proof that $\{B_m\}_{m \in \mathbb{N}}$ is a countable collection of nested, nonempty, closed subsets of A that has an empty intersection. \square

Exercise. Show that the $\{B_m\}_{m \in \mathbb{N}}$ constructed above is a countable collection of nested, nonempty, closed subsets of A that has an empty intersection.